arXiv:1509.01753v2 [math.AP] 26 Sep 2015

POSITIVE STEADY STATES OF AN INDEFINITE EQUATION WITH A NONLINEAR BOUNDARY CONDITION: EXISTENCE, MULTIPLICITY, STABILITY AND ASYMPTOTIC PROFILES

HUMBERTO RAMOS QUOIRIN AND KENICHIRO UMEZU

Abstract. We investigate positive steady states of an indefinite superlinear reaction-diffusion equation arising from population dynamics, coupled with a nonlinear boundary condition. Both the equation and the boundary condition depend upon a positive parameter λ, which is inversely proportional to the diffusion rate. We establish several multiplicity results when the diffusion rate is large and analyze the asymptotic profiles and the stability properties of these steady states as the diffusion rate grows to infinity. In particular, our results show that in some cases bifurcation from zero and from infinity occur at $\lambda = 0$. Our approach combines variational and bifurcation techniques.

1. Introduction and main results

Let Ω be a bounded and regular domain of $\mathbb{R}^N$ with $N \geq 2$. In this article we are concerned with the problem

$$(P_\lambda) \qquad \begin{cases} -\Delta u = \lambda(m(x)u + a(x)|u|^{p-2}u) & \text{in} \quad \Omega, \\ \frac{\partial u}{\partial \mathbf{n}} = \lambda b(x)|u|^{q-2}u & \text{on} \quad \partial\Omega, \end{cases}$$

where:

- Δ is the usual Laplacian in $\mathbb{R}^N$
- $\lambda \in \mathbb{R}$
- $1 < q < 2 < p$ and if $N > 2$ then $p < 2^* = \frac{2N}{N-2}$
- $m, a \in L^\infty(\Omega)$, $m^+ \not\equiv 0$
- $b \in L^\infty(\partial\Omega)$
- $\mathbf{n}$ is the outward unit normal to $\partial\Omega$.

Our main goal in this article is to carry on the study of (P_λ), which was addressed in [23] for the logistic case $a \leq 0$. By variational and bifurcation techniques, we established existence and multiplicity results for non-negative solutions of (P_λ). Moreover, the structure of the non-negative solutions set was also discussed. We intend now to deal with these issues in the case where a changes sign.

By a solution of (P_λ) we mean a weak solution, i.e. $u \in H^1(\Omega)$ satisfying

$$\int_\Omega \nabla u \nabla \varphi - \lambda \int_\Omega m(x) u\varphi - \lambda \int_\Omega a(x)|u|^{p-2}u\varphi - \lambda \int_{\partial\Omega} b(x)|u|^{q-2}u\varphi = 0, \quad \forall \varphi \in H^1(\Omega).$$

In this case, we may also say that the couple (λ, u) is a solution of (P_λ). As already pointed out in [23], solutions of (P_λ) satisfy $u \in W^{2,r}_{\mathrm{loc}}(\Omega) \cap \mathcal{C}^\theta(\overline{\Omega})$ for some $r > N$ and $0 < \theta < 1$, so

---

1991 *Mathematics Subject Classification.* 35J20, 35J25, 35J61, 35B32, 35P30.

Key words and phrases. Semilinear elliptic problem, Indefinite weight, Variational methods, Bifurcation approach, Population dynamics.

The first author was supported by the FONDECYT grant 11121567.

that by the weak maximum principle [17, Theorem 9.1], nontrivial non-negative solutions of (P_λ) are strictly positive in Ω.

(P_λ) describes the steady state of solutions of the corresponding initial boundary value problem

$$\begin{cases} \dfrac{\partial u}{\partial t} = \nabla \cdot (d\nabla u) + (m(x) + a(x)|u|^{p-2})u & \text{in } (0,\infty)\times\Omega, \\ u(0,x) = u_0(x) \geq 0 & \text{in } \Omega, \\ (d\nabla u)\cdot \mathbf{n} = b(x)|u|^{q-2}u & \text{on } (0,\infty)\times\partial\Omega, \end{cases} \tag{1.1}$$

which appears as a model in population dynamics (see Cantrell and Cosner [12], Gómez-Reñasco and López-Gómez [18]). Here the unknown function u stands for the population density of some species having $m(x)$ as intrinsic growth rate and $m(x) + a(x)|u|^{p-2}$ as extrinsic growth rate. If $a(x) \leq 0$ then the latter one is the well known logistic growth rate, with self-limitation ($a(x) < 0$) or without limitation ($a(x) = 0$), so that the region where $a(x) = 0$ can be considered as a *refuge*. We can give the case $a(x) > 0$ the following biological interpretation (cf. [18]): in this case, the extrinsic growth rate measures the *symbiosis* effect due to the intraspecific cooperation whereas, in the case $a(x) < 0$, it measures the crowding effect associated with competition. As for the nonlinear boundary condition, it suggests that the flux rate $(d\nabla u)\cdot\mathbf{n}$ of the population on $\partial\Omega$ is incoming or outgoing (according to the sign of $b(x)$) and depends nonlinearly on u as $|u|^{q-2}u$ (cf. [15]). From the population dynamics viewpoint, we point out that the parameter λ appearing in (P_λ) describes the reciprocal number of the diffusion coefficient $d > 0$, and only non-negative solutions are of interest.

Elliptic problems with indefinite nonlinearities have been studied over the last 25 years, starting with the works of Bandle, Pozio and Tesei [4], Ouyang [22], Alama and Tarantello [2], Berestycki, Capuzzo-Dolcetta and Nirenberg [5, 6], Lopez-Gomez [19], etc. Brown and Zhang [11] and Brown [8] used the Nehari manifold (or fibering) method to discuss existence, multiplicity, and non-existence of positive solutions for the problem

$$\begin{cases} -\Delta u = \lambda m(x)u + a(x)|u|^{p-2}u & \text{in} \quad \Omega, \\ u = 0 & \text{on} \quad \partial\Omega, \end{cases}$$

according to the position of λ. The sublinear case $1 < p < 2$ and the superlinear case $p > 2$ were treated in [8] and [11], respectively. We shall see in this article that the Nehari manifold method turns out to be efficient for (P_λ) as well.

We note that (P_λ) is characterized by the combination of the nonlinearities $m(x)u + a(x)|u|^{p-2}u$ in Ω and $b(x)|u|^{q-2}u$ on $\partial\Omega$. Furthermore, the signs of m, a and b may completely change the effect of these nonlinearities. For elliptic problems with such combined nonlinearities, we refer to Chipot, Fila, and Quittner [13], López-Gómez, Márquez, and Wolanski [20], Morales-Rodrigo and Suárez [21], Wu [31]. Besides investigating existence and multiplicity of positive solutions, we shall analyze as well the structure of the positive solutions set. Our approach is mainly based on a detailed study of the energy functional associated to (P_λ). Note however that unlike most of the aforementioned works, (P_λ) lacks coercivity on its left-hand side, since the term $(\int_\Omega |\nabla u|^2)^{\frac{1}{2}}$ does not correspond to the norm of u in $H^1(\Omega)$.

Since the set of solutions of (P_λ) at $\lambda = 0$ is explicitly provided by the constants, we may obtain positive solutions for $|\lambda|$ small by a bifurcation analysis on the line $(\lambda, u) = (0, c)$, where $c > 0$ is a constant, cf. [28, 29]. On the other hand, the lack of continuity of the derivative of $|u|^{q-2}u$ at $u = 0$ prevents the use of the bifurcation approach to obtain positive solutions bifurcating from the zero solution. Finally, let us remark that the boundary point lemma cannot be applied directly to our problem, since it seems difficult to deduce that any nontrivial non-negative solution belongs to $\mathcal{C}^1(\overline{\Omega})$ in view of the assumption $0 < q-1 < 1$. Therefore we are not able to infer that nontrivial non-negative solutions are positive on $\overline{\Omega}$. However we shall prove in Proposition 5.1 that if u is a nontrivial non-negative solution of (P_λ) then the set $\{x \in \partial\Omega : u(x) = 0\}$ has no interior points in the relative topology of $\partial\Omega$. Note that, in the one-dimensional case $N = 1$, the boundary point lemma is applicable, and we can deduce that any nontrivial non-negative weak solution of (P_λ) is positive on $\overline{\Omega}$, see Proposition B.1 in Appendix B.

Let us set the notations and conventions used in this article:

- The infimum of an empty set is assumed to be ∞.
- Unless otherwise stated, for any $f \in L^1(\Omega)$ the integral $\int_\Omega f$ is considered with respect to the Lebesgue measure, whereas for any $g \in L^1(\partial\Omega)$ the integral $\int_{\partial\Omega} g$ is considered with respect to the surface measure.
- For $r \geq 1$ the Lebesgue norm in $L^r(\Omega)$ will be denoted by $\|\cdot\|_r$ and the usual norm of $H^1(\Omega)$ by $\|\cdot\|$.
- We set $2^* = \frac{2N}{N-2}$ and $2_* = \frac{2(N-1)}{N-2}$ for $N > 2$.
- The strong and weak convergence are denoted by $\rightarrow$ and $\rightharpoonup$, respectively.
- The positive and negative parts of a function u are defined by $u^\pm := \max\{\pm u, 0\}$.
- If $U \subset \mathbb{R}^N$ then we denote the closure of U by $\overline{U}$ and the interior of U by $\operatorname{int} U$.
- The support of a measurable function f is denoted by $\operatorname{supp} f$.
- If Φ is a functional defined on $H^1(\Omega)$, we set

$$\Phi^\pm := \{u \in H^1(\Omega);\ \Phi(u) \gtrless 0\}, \quad \Phi_0 := \Phi^{-1}(0) \quad \text{and} \quad \Phi_0^\pm := \Phi^\pm \cup \Phi_0.$$

Recall that

$$\lambda_1 = \lambda_1(m) := \inf\left\{\int_\Omega |\nabla u|^2; u \in H^1(\Omega), \int_\Omega mu^2 = 1\right\} \tag{1.2}$$

is a principal and simple eigenvalue of the problem

$$\begin{cases} -\Delta u = \lambda m(x) u & \text{in} \quad \Omega, \\ \frac{\partial u}{\partial \mathbf{n}} = 0 & \text{on} \quad \partial\Omega. \end{cases}$$

It is well known (cf. Brown and Lin [10]) that $\lambda_1(m) > 0$ if and only if $\int_\Omega m < 0$, in which case $\lambda_1(m)$ is achieved by a unique non-constant eigenfunction φ_1 such that $\varphi_1 > 0$ on $\overline{\Omega}$. If $\int_\Omega m > 0$ then $\lambda_1(m) = 0$ is achieved by $\varphi_1 = \left(\int_\Omega m\right)^{-\frac{1}{2}}$.

We set

$$C_{pq} = \frac{q(p-2)}{2(p-q)} \left(\frac{p(2-q)}{2(p-q)}\right)^{\frac{2-q}{p-2}} \tag{1.3}$$

and

$$K_1(m,a) = C_{pq} \frac{\left(\int_\Omega m\right)^{\frac{p-q}{p-2}}}{\left(-\int_\Omega a\right)^{\frac{2-q}{p-2}}}, \tag{1.4}$$

whenever $\int_\Omega m > 0 > \int_\Omega a$.

Let $\varphi : [0, \infty) \rightarrow \mathbb{R}$ be given by

$$\varphi(t) = t^{2-q} \int_\Omega m + t^{p-q} \int_\Omega a + \int_{\partial\Omega} b. \tag{1.5}$$

It is easily seen that if either

$$\int_\Omega m > 0 > \int_\Omega a \quad \text{and} \quad -K_1(m,a) < \int_{\partial\Omega} b < 0$$

or

$$\int_\Omega m < 0 < \int_\Omega a \quad \text{and} \quad K_1(-m,-a) > \int_{\partial\Omega} b > 0$$

then φ has two zeros $c_1 < c_2$, which satisfy

$$c_1 < \left(-\frac{(2-q)\int_\Omega m}{(p-q)\int_\Omega a}\right)^{\frac{1}{p-2}} < c_2. \tag{1.6}$$

If either

$$\int_\Omega a < 0 < \int_{\partial\Omega} b \quad \text{or} \quad \int_\Omega a > 0 > \int_{\partial\Omega} b$$

then φ has an unique zero, denoted by c_0.

In order to simplify the notation, we introduce the functionals

$$E_\lambda(u) = \int_\Omega \left(|\nabla u|^2 - \lambda m(x) u^2\right), \quad A(u) = \int_\Omega a(x)|u|^p \quad \text{and} \quad B(u) = \int_{\partial\Omega} b(x)|u|^q,$$

defined on $H^1(\Omega)$. From the compactness of the Sobolev embeddings

$$H^1(\Omega) \hookrightarrow L^r(\Omega), \quad H^1(\Omega) \hookrightarrow L^t(\partial\Omega),$$

for $r \in [1, 2^*)$ and $t \in [1, 2_*)$ respectively, it is straightforward that E_λ is weakly lower semi-continuous, whereas A and B are weakly continuous.

Remark 1.1. The following results, which can be easily verified, will be used repeatedly throughout this article:

(1) If (u_n) is a bounded sequence in $H^1(\Omega)$ then we may assume that $u_n \rightharpoonup u_0$, $A(u_n) \to A(u_0)$, $B(u_n) \to B(u_0)$, and $E_\lambda(u_0) \le \liminf E_\lambda(u_n)$, for some $u_0 \in H^1(\Omega)$.

(2) If $u_n \rightharpoonup 0$ in $H^1(\Omega)$ and $\limsup E_\lambda(u_n) \le 0$ then $u_n \to 0$ in $H^1(\Omega)$.

Let us set

$$\lambda_a = \lambda_a(m) := \inf\left\{\int_\Omega |\nabla u|^2; u \in A_0^+, \int_\Omega m u^2 = 1\right\}, \tag{1.7}$$

$$\lambda_b = \lambda_b(m) := \inf\left\{\int_\Omega |\nabla u|^2; u \in B_0^+, \int_\Omega m u^2 = 1\right\}, \tag{1.8}$$

and

$$\lambda_s = \lambda_s(m,a,b) = \inf\left\{\int_\Omega |\nabla u|^2;\ u \in A_0^+ \cap B_0^+,\ S(u) = 1\right\}, \tag{1.9}$$

where

$$S(u) = \int_\Omega m u^2 + \left(C_{pq}^{-1} B(u) A(u)^{\frac{2-q}{p-2}}\right)^{\frac{p-2}{p-q}} \tag{1.10}$$

is defined for $u \in A_0^+ \cap B_0^+$. One may easily check that S is quadratic, i.e. $S(tu) = t^2 S(u)$ for $t \in \mathbb{R}$ and $u \in A_0^+ \cap B_0^+$. Lastly, we set

$$\Omega_\pm := \text{int supp } a^\pm. \tag{1.11}$$

We are now in position to state our main results. To begin with, we state an existence result, which provides bifurcation from zero and from infinity at $\lambda = 0$ in the following sense:

Definition 1.2. It is said that *bifurcation from zero occurs at* $\lambda = \lambda_0 \in \mathbb{R}$ for (P_λ) if there exist nontrivial nonnegative solutions u_j of (P_{λ_j}) such that $\lambda_j \to \lambda_0$, and $u_j \to 0$ in $\mathcal{C}(\overline{\Omega})$ as $j \to \infty$. Similarly, it is said that *bifurcation from infinity occurs at* $\lambda = \lambda_\infty \in \mathbb{R}$ for (P_λ) if there exist nontrivial nonnegative solutions u_j of (P_{λ_j}) such that $\lambda_j \to \lambda_\infty$, and $\|u_j\|_{\mathcal{C}(\overline{\Omega})} \to \infty$ as $j \to \infty$.

Theorem 1.3.

1. *If $\int_{\partial\Omega} b < 0$ then $\lambda_b, \lambda_s > 0$. If, in addition, $b^+ \not\equiv 0$ then (P_λ) has a nontrivial non-negative solution $u_{0,\lambda}$ for $0 < \lambda < \min\{\lambda_b, \lambda_s\}$, which satisfies $u_{0,\lambda} \to 0$ in $\mathcal{C}^\theta(\overline{\Omega})$ for some $\theta \in (0,1)$ as $\lambda \to 0^+$. Moreover, there exists $\lambda_n \to 0^+$ such that $\lambda_n^{-\frac{1}{2-q}} u_{0,\lambda_n} \to w_0$ in $H^1(\Omega) \cap \mathcal{C}^\theta(\overline{\Omega})$ as $n \to \infty$, where w_0 is a nontrivial non-negative solution of*
$$-\Delta w = 0 \quad \text{in } \Omega, \qquad \frac{\partial w}{\partial \mathbf{n}} = b(x) w^{q-1} \quad \text{on } \partial\Omega. \tag{1.12}$$
Furthermore, $w_0 > 0$ in Ω, the set $\{x \in \partial\Omega : w_0 = 0\}$ has no interior points in the relative topology of $\partial\Omega$, and it is contained in $\{x \in \partial\Omega : b(x) \le 0\}$ if $b \in \mathcal{C}(\partial\Omega)$.

2. *If $\int_\Omega a < 0$ then $\lambda_a, \lambda_s > 0$. If, in addition, $a^+ \not\equiv 0$ then (P_λ) has a nontrivial non-negative solution $u_{2,\lambda}$ for $0 < \lambda < \min\{\lambda_a, \lambda_s\}$, which satisfies $u_{2,\lambda} > 0$ in $\overline{\Omega}$ for $\lambda > 0$ sufficiently small and $\min_{\overline{\Omega}} u_{2,\lambda} \to \infty$ as $\lambda \to 0^+$. Moreover, there exists $\lambda_n \to 0^+$ such that $\lambda_n^{\frac{1}{p-2}} u_{2,\lambda_n} \to w_\infty$ in $H^1(\Omega) \cap C^\theta(\overline{\Omega})$ for some $\theta \in (0,1)$ as $n \to \infty$, where $w_\infty \in W^{2,r}(\Omega)$, $r > N$, is a solution of the problem*
$$-\Delta w = a(x) w^{p-1} \quad \text{in } \Omega, \qquad \frac{\partial w}{\partial \mathbf{n}} = 0 \quad \text{on } \partial\Omega.$$
Furthermore $w_\infty > 0$ on $\overline{\Omega}$.

Remark 1.4.

1. Theorem 1.3 states that if $b^+ \not\equiv 0$ and $\int_{\partial\Omega} b < 0$ then (P_λ) has, besides the trivial branch $\{(\lambda, u) = (0, c); c \text{ is a positive constant}\}$, the solution $u_{0,\lambda}$ bifurcating from zero at $\lambda = 0$. In a similar way, if $a^+ \not\equiv 0$ and $\int_\Omega a < 0$ then (P_λ) has, besides the trivial branch of positive constants, the solution $u_{2,\lambda}$ bifurcating from infinity at $\lambda = 0$. Furthermore, the blow up of $u_{2,\lambda}$ as $\lambda \to 0^+$ occurs uniformly on $\overline{\Omega}$.

2. We shall see in Proposition 5.1 that for any $x \in \partial\Omega$ such that $b(x) > 0$ there holds $u_{0,\lambda}(x) > 0$ if $\lambda > 0$ is sufficiently small. More precisely, if $\Sigma \subset \partial\Omega$ is such that $\overline{\Sigma} \subset \{x \in \partial\Omega : b(x) > 0\}$ then $\inf_\Sigma u_{0,\lambda} > 0$ for $\lambda > 0$ sufficiently small.

Our next results provide multiplicity of non-negative solutions and are stated under the condition
$$\int_\Omega m > 0. \tag{1.13}$$
Since we are considering $\lambda \in \mathbb{R}$, the case $\int_\Omega m < 0$ reduces to (1.13) after the change of variable $\lambda \mapsto -\lambda$.

Theorem 1.5. *Assume* (1.13), *$\int_\Omega a < 0$, and $-K_1(m,a) < \int_{\partial\Omega} b < 0$. Then:*

1. *(P_λ) has a nontrivial non-negative solution $u_{1,\lambda}$ for $0 < \lambda < \min\{\lambda_a, \lambda_b\}$. Moreover $u_{1,\lambda} \to c_2$ in $\mathcal{C}^\theta(\overline{\Omega})$ for some $\theta \in (0,1)$ as $\lambda \to 0^+$.*

2. *(P_λ) has two nontrivial non-negative solutions $v_{1,\lambda}$, $v_{2,\lambda}$ for $\max\{\tilde{\lambda}_1, \tilde{\lambda}_s\} < \lambda < 0$, where $\tilde{\lambda}_1 = -\lambda_1(-m)$ and $\tilde{\lambda}_s = -\lambda_s(-m, -a, -b)$. Furthermore, $v_{1,\lambda} \to c_1$ and $v_{2,\lambda} \to c_2$ in $\mathcal{C}^\theta(\overline{\Omega})$ for some $\theta \in (0,1)$ as $\lambda \to 0^-$.*

Under the assumptions of Theorem 1.5 and stronger regularity conditions on m, a, b, we shall obtain, for $\lambda > 0$ sufficiently small, a positive solution of (P_λ) converging to c_1 as $\lambda \to 0^+$. This will be carried out by a bifurcation argument and, as a consequence, will provide at least four nontrivial non-negative solutions of (P_λ) for $\lambda > 0$ small enough. Since this argument requires only the existence of zeros of φ and some regularity on m, a, and b, we shall assume, in addition to (1.13),

$$m, a \in \mathcal{C}^\alpha(\overline{\Omega}) \quad \text{and} \quad b \in \mathcal{C}^{1+\alpha}(\partial\Omega), \quad \text{for some} \quad 0 < \alpha < 1, \tag{1.14}$$

and

$$\int_\Omega a < 0 \quad \text{and} \quad -\tilde{K}_1(m,a) < \int_{\partial\Omega} b < 0, \tag{1.15}$$

where

$$\tilde{K}_1(m,a) = \tilde{C}_{pq} \frac{\left(\int_\Omega m\right)^{\frac{p-q}{p-2}}}{\left(-\int_\Omega a\right)^{\frac{2-q}{p-2}}}, \tag{1.16}$$

and

$$\tilde{C}_{pq} = \left(\frac{q}{2} \left(\frac{p}{2}\right)^{\frac{2-q}{p-2}} \right)^{-1} C_{pq}.$$

It is easily seen that $\tilde{C}_{pq} > C_{pq}$, so that $\tilde{K}_1(m,a) > K_1(m,a)$.

Hereafter, by a classical positive solution of (P_λ) we mean $u \in \mathcal{C}^{2+\theta}(\overline{\Omega})$ for some $0 < \theta < 1$ which satisfies (P_λ) in the classical sense and is strictly positive on $\overline{\Omega}$.

Theorem 1.6. *Assume* (1.14). *If a changes sign and $\int_\Omega a < 0$, then, for $\lambda > 0$ sufficiently small, the solution $u_{2,\lambda}$ obtained in Theorem 1.3 is a classical positive solution of (P_λ), which is moreover unstable.*

Remark 1.7. Since $1 < q < 2$, solutions of (1.12) are in $\mathcal{C}^\theta(\overline{\Omega})$ with $\theta \in (0,1)$, so that the boundary point lemma is not applicable, and consequently we can not deduce the positivity of w_0 on $\partial\Omega$. If this is the case then we can prove a result similar to Theorem 1.6 for $u_{0,\lambda}$, obtained in Theorem 1.3. More precisely, let us assume (1.14) and $w_0 > 0$ on $\overline{\Omega}$. If b changes sign and $\int_{\partial\Omega} b < 0$ then, for $\lambda > 0$ sufficiently small, $u_{0,\lambda}$ is a classical positive solution of (P_λ) which is moreover asymptotically stable. We include a sketch of the proof of this result in Section 5.

Theorem 1.8. *Assume* (1.13), (1.14) *and* (1.15). *Then there exists $\underline{\lambda} > 0$ such that:*

(1) $u_{1,\lambda}$ is a classical positive solution of (P_λ) which is asymptotically stable for $\lambda \in (0, \underline{\lambda})$. Moreover, (P_λ) has a classical positive solution $u_{3,\lambda}$ for $\lambda \in (0, \underline{\lambda})$ which is unstable. These solutions are continuous in $\mathcal{C}^{2+\alpha}(\overline{\Omega})$ with respect to λ and emanate from $(0, c_2)$ and $(0, c_1)$ respectively, i.e. $u_{1,0} = c_2$ and $u_{3,0} = c_1$. Finally, there are no other classical positive solutions of (P_λ) converging to some positive constant in $\mathcal{C}(\overline{\Omega})$ as $\lambda \to 0^+$.

(2) $v_{1,\lambda}, v_{2,\lambda}$ are classical positive solutions of (P_λ) for $\lambda \in (-\underline{\lambda}, 0)$, which are asymptotically stable, and unstable, respectively. Moreover, these solutions are continuous in $\mathcal{C}^{2+\alpha}(\overline{\Omega})$ with respect to λ and emanate from $(0, c_1)$ and $(0, c_2)$ respectively, i.e. $v_{1,0} = c_1$ and $v_{2,0} = c_2$. Finally, there are no other classical positive solutions of (P_λ) converging to some positive constant in $\mathcal{C}(\overline{\Omega})$ as $\lambda \to 0^-$.

Remark 1.9. Theorem 1.8 states, in particular, that $v_{1,\lambda}$, $v_{2,\lambda}$ are the extensions of $u_{3,\lambda}$, $u_{1,\lambda}$, respectively, to the region $\lambda < 0$ (see Figure 1).

Combining Theorems 1.3 and 1.5, we get a multiplicity result for (P_λ) with $\lambda > 0$ small, which is obtained by variational techniques:

Corollary 1.10. *Under the assumptions of Theorem 1.5, assume moreover that* $a^+ \not\equiv 0$ *and* $b^+ \not\equiv 0$. *Then* (P_λ) *has three nontrivial non-negative solutions* $u_{j,\lambda}$, $j = 0, 1, 2$, *for* $0 < \lambda < \min\{\lambda_s, \lambda_a, \lambda_b\}$, *which satisfy* $u_{0,\lambda} \to 0$ *and* $u_{1,\lambda} \to c_2$ *in* $\mathcal{C}^\theta(\overline{\Omega})$ *for some* $\theta \in (0,1)$ *and* $\min\limits_{\overline{\Omega}} u_{2,\lambda} \to \infty$ *as* $\lambda \to 0^+$.

From Corollary 1.10 and Theorem 1.8(1), we infer the following multiplicity result, which is obtained combining the variational and bifurcation approaches:

Corollary 1.11. *In addition to the assumptions of Theorem 1.5, we assume* (1.14). *Assume moreover* $a^+ \not\equiv 0$. *Then* (P_λ) *has at least three classical positive solutions for* $0 < \lambda < \underline{\lambda}$, *namely* $u_{j,\lambda}$, $j = 1, 2, 3$. *If, in addition,* $b^+ \not\equiv 0$ *then* (P_λ) *has a fourth nontrivial non-negative solution for* $0 < \lambda < \underline{\lambda}$, *namely,* $u_{0,\lambda}$.

Remark 1.12. We shall see in Proposition 5.3 that under (1.13), (1.14), and (1.15), (P_λ) has, for $|\lambda|$ sufficiently small, two classical positive solutions $U_{j,\lambda}$, $j = 1, 2$, which are continuous (with respect to λ) in $\mathcal{C}^{2+\alpha}(\overline{\Omega})$ and satisfy $U_{j,0} = c_j$, $j = 1, 2$. Furthermore, this result does not require the condition $p < 2^*$. So, under (1.13), (1.14) and the conditions

$$a, b \text{ change sign}, \quad \int_\Omega a < 0, \quad \text{and} \quad \int_{\partial\Omega} b < 0,$$

Theorem 1.3 and Proposition 5.3 provide, according to the value of $\int_{\partial\Omega} b$, the following result on the number of nontrivial non-negative solutions of (P_λ):

(1) If $\int_{\partial\Omega} b > -\tilde{K}_1(m, a)$ then (P_λ) has at least four nontrivial non-negative solutions (two variational nontrivial non-negative solutions, among which one is a classical positive solution, and two bifurcating classical positive solutions) for $\lambda > 0$ sufficiently small.

(2) If $\int_{\partial\Omega} b \le -\tilde{K}_1(m, a)$ then (P_λ) has at least two (variational) nontrivial non-negative solutions, among which one is a classical positive solution, for $\lambda > 0$ sufficiently small. Moreover, if $\int_{\partial\Omega} b < -\tilde{K}_1(m, a)$ then (P_λ) has no classical positive solutions converging to a positive constant as $\lambda \to 0^+$.

The following condition provides an *a priori* bound on the values of λ for which (P_λ) has a nontrivial non-negative solution:

$$(\mathcal{H}) \qquad \begin{cases} \text{There are smooth sub-domains } D_\pm \Subset \Omega \text{ such that} \\ D_\pm \subset \Omega_\pm \text{ and } m \text{ changes sign in } D_+ \text{ and in } D_-. \end{cases}$$

Theorem 1.13. *Assume that* a *changes sign and* $(\mathcal{H})$ *holds. Then there exists a constant* $\Lambda > 0$ *such that if* u *is a nontrivial non-negative solution of* (P_λ) *then* $|\lambda| \le \Lambda$.

As a particular case of (P_λ), we shall consider the problem

$$(Q_\lambda) \qquad \begin{cases} -\Delta u = \lambda m(x)(u - |u|^{p-2}u) & \text{in} \quad \Omega, \\ \frac{\partial u}{\partial \mathbf{n}} = \lambda b(x)|u|^{q-2}u & \text{on} \quad \partial\Omega, \end{cases}$$

where λ, m, b, p, q are as above. Note that (Q_λ) corresponds to (P_λ) with $a \equiv -m$. The nonlinearity $\lambda m(x)(u - |u|^{p-2}u)$ arises from population genetics and has already been studied under homogeneous boundary conditions in [9, 14, 27].

Let

$$\lambda_r = \lambda_r(m,b) := \inf\left\{\int_\Omega |\nabla u|^2;\ u\in H^1(\Omega),\ \int_\Omega mu^2=1, \int_\Omega m|u|^p \le 0 \le \int_{\partial\Omega} b|u|^q\right\},$$

$$\lambda_t = \lambda_t(m,b) := \inf\left\{\int_\Omega |\nabla u|^2;\ u\in H^1(\Omega),\ \int_\Omega m|u|^p \le 0 \le \int_{\partial\Omega} b|u|^q,\ T(u)=1\right\},$$

where

$$T(u) = \int_\Omega mu^2 + \left(C_{pq}^{-1}\left(\int_{\partial\Omega} b|u|^q\right)\left(-\int_\Omega m|u|^p\right)^{\frac{2-q}{p-2}}\right)^{\frac{p-2}{p-q}},$$

is defined for $u\in H^1(\Omega)$ such that $\int_\Omega m|u|^p \le 0 \le \int_{\partial\Omega} b|u|^q$. Note that $\lambda_r = \min\{\lambda_a(m), \lambda_b(m)\}$ and $\lambda_t = \lambda_s(m,a,b)$ with $a\equiv -m$.

Finally, let

$$\phi(t) = \left(t^{2-q}-t^{p-q}\right)\int_\Omega m + \int_{\partial\Omega} b.$$

The assumption of Theorem 1.5 reads now

$$\int_\Omega m > 0 > \int_{\partial\Omega} b > -C_{pq}\int_\Omega m, \tag{1.17}$$

in which case ϕ has two positive zeros $c_1 < c_2$.

Applying Theorems 1.3 and 1.5 to (Q_λ) we obtain the following:

Theorem 1.14. *Assume* (1.17) *and m changes sign. Then $\lambda_r, \lambda_t > 0$ and:*

(1) (Q_λ) has two nontrivial non-negative solutions $u_{1,\lambda}$, $u_{2,\lambda}$ for $0<\lambda<\min\{\lambda_1,\lambda_t\}$, which satisfy $u_{1,\lambda}\to c_2$ in $\mathcal{C}^\theta(\overline{\Omega})$ for some $\theta\in(0,1)$ and $\min_{\overline{\Omega}} u_{2,\lambda}\to\infty$ as $\lambda\to 0^+$.

If, in addition, $b^+\not\equiv 0$, then (Q_λ) has a further nontrivial non-negative solution $u_{0,\lambda}$ for $0<\lambda<\lambda_r$, which satisfies $u_{0,\lambda}\to 0$ in $\mathcal{C}^\theta(\overline{\Omega})$ as $\lambda\to 0^+$.

(2) (Q_λ) has two nontrivial non-negative solutions $v_{1,\lambda}$, $v_{2,\lambda}$ for $\max\{\tilde\lambda_1,\tilde\lambda_t\}<\lambda<0$, where $\tilde\lambda_1 = -\lambda_1(-m)$ and $\tilde\lambda_t = -\lambda_t(-m,-b)$. Furthermore, $v_{1,\lambda}\to c_1$ and $v_{2,\lambda}\to c_2$ in $\mathcal{C}^\theta(\overline{\Omega})$ for some $\theta\in(0,1)$ as $\lambda\to 0^-$.

Lastly, we consider the case where ϕ has a unique positive zero c_0 such that $\phi'(c_0)=0$, which occurs precisely when

$$\int_\Omega m > 0 > \int_{\partial\Omega} b = -\tilde K_1(m,-m), \tag{1.18}$$

where $\tilde K_1$ is defined in (1.16). Under (1.14), we obtain a smooth curve of classical positive solutions of (Q_λ) for $|\lambda|$ sufficiently small by a bifurcation argument.

The following result asserts that if (1.18) holds then $(0,c_0)$ is a turning point to the right on the smooth curve of positive solutions of (Q_λ).

Theorem 1.15. *Assume* (1.14). *If (Q_λ) has a classical positive solution u with $\lambda\neq 0$ such that $u_\lambda\to c$ in $\mathcal{C}(\overline{\Omega})$ as $\lambda\to 0$, where c is a positive constant, then $\phi(c)=0$. Moreover:*

(1) Assume (1.15) *with $a=-m$. Then there exist two arbitrarily smooth maps $\lambda\mapsto U_{1,\lambda}, U_{2,\lambda}\in\mathcal{C}^{2+\alpha}(\overline{\Omega})$ for λ close to 0 such that $U_{1,\lambda}$, $U_{2,\lambda}$ are classical positive solutions of (Q_λ) and satisfy $U_{1,0}=c_1$ and $U_{2,0}=c_2$.*

(2) *Assume* (1.18). *Then there exist two arbitrarily smooth maps* $t \mapsto \lambda(t) \in \mathbb{R}$ *and* $t \mapsto u(t) \in \mathcal{C}^{2+\alpha}(\overline{\Omega})$ *for* t *close to* c_0 *such that* $(\lambda, u) = (\lambda(t), u(t))$ *is a classical positive solution of* (Q_λ) *with* $\lambda(c_0) = \lambda'(c_0) = 0$*,* $\lambda''(c_0) > 0$*, and* $u(t) = t + o(1)$ *as* $t \to c_0$*. Moreover, there exists a constant* $\varepsilon > 0$ *such that* $u(t)$ *is asymptotically stable for* $c_0 < t < c_0 + \varepsilon$ *and unstable for* $c_0 - \varepsilon < t < c_0$*.*

Remark 1.16.

(1) Theorem 1.15 holds in fact without the condition $p < 2^*$.
(2) Assertion (1) in Theorem 1.15 corresponds to Theorem 1.8 for (Q_λ).

Remark 1.17. The results on the number of nontrivial non-negative solutions from Remark 1.12 can be sharpened for (Q_λ). In this case, Theorems 1.3 and 1.15 provide that, under (1.14), if m and b change sign, and $\int_\Omega m > 0 > \int_{\partial\Omega} b$, then (Q_λ) has:

(1) at least four nontrivial non-negative solutions (two variational nontrivial non-negative solutions, among which one is a classical positive solution, and two bifurcating classical positive solutions) for $\lambda > 0$ sufficiently small if $\int_{\partial\Omega} b \geq -\tilde{K}_1(m, -m)$.

(2) at least two (variational) nontrivial non-negative solutions, among which one is a classical positive solution, for $\lambda > 0$ sufficiently small and no classical positive solutions converging to a positive constant as $\lambda \to 0^+$ if $\int_{\partial\Omega} b < -\tilde{K}_1(m, -m)$.

1.1. Suggested bifurcation diagrams.

In view of the results stated above, we analyze now the possible bifurcations diagrams of (P_λ) and (Q_λ). This will be done assuming that a changes sign and $(\mathcal{H})$ holds, in which case Theorem 1.13 ensures that (P_λ) has no nontrivial non-negative solution for $|\lambda| > \Lambda$.

(1) Assume (1.13), a and b change sign, and $\int_\Omega a < 0$. If $-K_1(m, a) < \int_{\partial\Omega} b < 0$ then, by Theorems 1.3, 1.5, and 1.8, the bifurcation diagram for (P_λ) is suggested by Figure 1. Moreover, Remark 1.12 suggests that this bifurcation diagram approaches the one shown in Figure 2 as $\int_{\partial\Omega} b \to -\infty$. Indeed, note that the value Λ provided by Theorem 1.13 does not depend on b. Furthermore, it can be shown that λ_b and λ_s stay bounded away from zero if b^+ is bounded from above (cf. Remark 3.5). By a formal observation, the nonlinear boundary condition in (P_λ) approaches the Dirichlet boundary condition $u = 0$ as $\int_{\partial\Omega} b \to -\infty$. So, after the change of variable $v = \lambda^{\frac{1}{p-2}} u$ for $u \geq 0$, the limiting problem for (P_λ) when $\int_{\partial\Omega} b \to -\infty$ would be
$$-\Delta v = \lambda m(x) v + a(x) v^{p-1} \text{ in } \Omega, \quad v = 0 \text{ on } \partial\Omega.$$
This problem has been investigated by Ouyang [22] in the case $m \equiv 1$ and the existence of a single turning point in the positive solutions set has been proved under some conditions on a. We expect then that the bifurcation diagram of (P_λ) has a unique turning point if $\int_{\partial\Omega} b \ll 0$.

(2) Assume (1.13), $b \leq 0$, a changes sign, and $\int_\Omega a < 0$.

   (a) If $-K_1(m, a) < \int_{\partial\Omega} b < 0$ then the bifurcation diagram for (P_λ) in the case $b \leq 0$ is suggested by Figure 3. This is motivated by Theorems 1.3, 1.5, and 1.8, and the fact that $u_{0,\lambda}$ approaches zero as b^+ converges to zero. More precisely, in Proposition 3.6 we prove that if $b_n \to b$ in $L^\infty(\partial\Omega)$ with $b_n^+ \not\equiv 0$ for every n, then there exists $\lambda_0 > 0$ such that the solution u_{0,λ,b_n} exists for every n, and satisfies $u_{0,\lambda,b_n} \to 0$ in $\mathcal{C}^\theta(\overline{\Omega})$ for some $\theta \in (0, 1)$ and $0 < \lambda < \lambda_0$. Note that in

[24] we have obtained bifurcation from zero in the case $b \leq 0$ and $b \not\equiv 0$ when $a < 0$.

(b) In a similar way, if $-\tilde{K}_1(m,a) - \varepsilon < \int_{\partial\Omega} b < -\tilde{K}_1(m,a)$ with $\varepsilon > 0$ sufficiently small then the bifurcation diagram for (P_λ) would be suggested by Figure 4. Lastly, as $\int_{\partial\Omega} b \to -\infty$, the bifurcation diagram for (P_λ) is expected to approach Figure 5.

(3) Assume that m and b change sign, and $\int_\Omega m > 0 > \int_{\partial\Omega} b = -\tilde{K}_1(m,-m)$. By Remark 1.17, the bifurcation diagram of (Q_λ) is suggested by Figure 6.

Remark 1.18. From the biological viewpoint, the bifurcation diagram suggested in Figure 1 would provide three possible conditional states for the population as the diffusion rate grows to infinity, namely: extinction, explosion and persistence with a spatially uniform distribution.



FIGURE 1. Suggested bifurcation diagram for (P_λ) when a and b change sign, $\int_\Omega m > 0 > \int_\Omega a$, and $0 > \int_{\partial\Omega} b > -K_1(m,a)$.



FIGURE 2. Suggested bifurcation diagram for (P_λ) when a and b change sign, $\int_\Omega m > 0 > \int_\Omega a$, and $\int_{\partial\Omega} b \to -\infty$.



FIGURE 3. Suggested bifurcation diagram for (P_λ) when a changes sign, $b \leq 0$, $\int_\Omega m > 0 > \int_\Omega a$, and $0 > \int_{\partial\Omega} b > -K_1(m,a)$.



FIGURE 4. Suggested bifurcation diagram for (P_λ) when a changes sign, $b \leq 0$, $\int_\Omega m > 0 > \int_\Omega a$, and $\int_{\partial\Omega} b < -\tilde{K}_1(m,a)$.



(a) Bifurcation diagram with a single turning point.

(b) Bifurcation diagram without a turning point.

FIGURE 5. Suggested bifurcation diagram for (P_λ) when a changes sign, $b \leq 0$, $\int_\Omega m > 0 > \int_\Omega a$, and $\int_{\partial\Omega} b \to -\infty$.



FIGURE 6. Suggested bifurcation diagram for (Q_λ) when m and b change sign and $\int_\Omega m > 0 > \int_{\partial\Omega} b = -\tilde{K}_1(m, -m)$.

The rest of this article is organized as follows: in Section 2 we collect some preliminary results on λ_a, λ_b, λ_s and $\mathcal{N}_\lambda$, the Nehari manifold associated to (P_λ). In Section 3, we obtain two minimizers for I_λ (the energy functional associated to (P_λ)) constrained to $\mathcal{N}_\lambda$. These minimizers are located in $\mathcal{N}_\lambda^+$ and correspond to local minimizers of I_λ. In Section 4, we obtain a further minimizer of I_λ constrained to $\mathcal{N}_\lambda^-$, which corresponds to a minimax critical point of I_λ. Finally, in Section 5, we carry out a bifurcation analysis of (P_λ) and prove the theorems stated above.

2. PRELIMINARIES

Solutions of (P_λ) shall be obtained as critical points of the functional

$$I_\lambda(u) = \frac{1}{2}E_\lambda(u) - \frac{\lambda}{p}A(u) - \frac{\lambda}{q}B(u),$$

defined on $H^1(\Omega)$. It is straightforward that I_λ is weakly lower semicontinuous for any $\lambda \in \mathbb{R}$. In contrast with the case $a < 0$, for any value of λ the functional I_λ is not coercive. Consequently we shall restrict I_λ to its associated Nehari manifold, which is given by

$$\begin{aligned} \mathcal{N}_\lambda &= \{u \in H^1(\Omega) \setminus \{0\};\, \langle I'_\lambda(u), u\rangle = 0\} \\ &= \left\{u \in H^1(\Omega) \setminus \{0\};\, E_\lambda(u) = \lambda\,[A(u) + B(u)]\right\}. \end{aligned}$$

$\mathcal{N}_\lambda$ is naturally split into $\mathcal{N}_\lambda^+$, $\mathcal{N}_\lambda^-$ and $\mathcal{N}_\lambda^0$, given by

$$\mathcal{N}_\lambda^\pm = \{u \in \mathcal{N}_\lambda;\, \langle J'_\lambda(u), u\rangle \gtrless 0\} \quad \text{and} \quad \mathcal{N}_\lambda^0 = \{u \in \mathcal{N}_\lambda;\, \langle J'_\lambda(u), u\rangle = 0\},$$

where $J_\lambda(u) = \langle I'_\lambda(u), u\rangle$ for $u \in H^1(\Omega)$. Thus

$$\mathcal{N}_\lambda^\pm = \left\{u \in \mathcal{N}_\lambda;\, E_\lambda(u) \lesseqgtr \lambda\frac{p-q}{p-2}B(u)\right\} = \left\{u \in \mathcal{N}_\lambda;\, E_\lambda(u) \gtreqless \lambda\frac{p-q}{2-q}A(u)\right\}.$$

and

$$\mathcal{N}_\lambda^0 = \left\{u \in \mathcal{N}_\lambda;\, E_\lambda(u) = \lambda\frac{p-q}{p-2}B(u) = \lambda\frac{p-q}{2-q}A(u)\right\}.$$

It is well-known that $\mathcal{N}_\lambda \setminus \mathcal{N}_\lambda^0$ is a $\mathcal{C}^1$ manifold defined by a natural constraint, i.e. any critical point of the restriction of I_λ to $\mathcal{N}_\lambda \setminus \mathcal{N}_\lambda^0$ is a critical point of I_λ (see for instance [11, Theorem 2.3]). Given $u \in H^1(\Omega) \setminus \{0\}$, we set

$$j_u(t) = I_\lambda(tu), \quad t > 0.$$

Then $tu \in \mathcal{N}_\lambda^\pm$ if, and only if, $j_u'(t) = 0$ and $j_u''(t) \gtrless 0$. We shall use the map j_u to deduce some properties of $\mathcal{N}_\lambda$. Note that $j_u(t) = t^q i_u(t)$, where

$$i_u(t) = \frac{1}{2}t^{2-q}E_\lambda(u) - \frac{\lambda}{p}t^{p-q}A(u) - \frac{\lambda}{q}B(u) \tag{2.1}$$

for $t > 0$.

Remark 2.1. It is easily seen that c is a positive zero of φ if and only if $c \in \mathcal{N}_\lambda$, for any $\lambda \in \mathbb{R}$. Additionally, if $\varphi'(c) \gtrless 0$ then $c \in \mathcal{N}_\lambda^\pm$. More precisely:

1. If $\int_\Omega m > 0 > \int_\Omega a$ and $-K_1(m,a) < \int_{\partial\Omega} b < 0$ then $c_1 \in \mathcal{N}_\lambda^-$ and $c_2 \in \mathcal{N}_\lambda^+$.
2. If $\int_\Omega m < 0 < \int_\Omega a$ and $0 < \int_{\partial\Omega} b < K_1(-m,-a)$ then $c_1 \in \mathcal{N}_\lambda^+$ and $c_2 \in \mathcal{N}_\lambda^-$.
3. If $\int_\Omega a > 0 > \int_{\partial\Omega} b$ then $c_0 \in \mathcal{N}_\lambda^-$.
4. If $\int_\Omega a < 0 < \int_{\partial\Omega} b$ then $c_0 \in \mathcal{N}_\lambda^+$.

The role of λ_s in the study of $\mathcal{N}_\lambda$ becomes clear in the next result:

Proposition 2.2. *If either* $\int_\Omega a < 0$ *or* $\int_{\partial\Omega} b < 0$ *or*

$$\int_\Omega m < 0 < \int_\Omega a \quad \textit{and} \quad 0 < \int_{\partial\Omega} b < K_1(-m,-a). \tag{2.2}$$

then λ_s*, given by* (1.9)*, is achieved. In particular,* $\lambda_s > 0$*. Furthermore, if* $\lambda \in (0, \lambda_s)$ *then for every* $u \in B^+ \cap E_\lambda^+ \cap A^+$ *there are* $0 < t_1 < t_2$ *such that* $t_1 u \in \mathcal{N}_\lambda^+$ *and* $t_2 u \in \mathcal{N}_\lambda^-$*, and there is no other* $t > 0$ *such that* $tu \in \mathcal{N}_\lambda$*.*

Proof. Let (u_n) be a minimizing sequence for λ_s, i.e.

$$A(u_n) \geq 0, \quad B(u_n) \geq 0, \quad S(u_n) = 1 \quad \text{and} \quad \int_\Omega |\nabla u_n|^2 \to \lambda_s.$$

If (u_n) is unbounded then we set $v_n = \frac{u_n}{\|u_n\|}$. We may assume that

$$v_n \rightharpoonup v_0, \quad A(v_n) \to A(v_0), \quad \text{and} \quad B(v_n) \to B(v_0).$$

Thus

$$A(v_0) \geq 0, \quad B(v_0) \geq 0, \quad S(v_0) = 0, \quad \text{and} \quad \int_\Omega |\nabla v_n|^2 \to 0.$$

Hence v_0 is a nonzero constant, so that

$$\int_\Omega a \geq 0, \quad \int_{\partial\Omega} b \geq 0 \quad \text{and} \quad \int_\Omega m + \left(C_{pq}^{-1} \left(\int_{\partial\Omega} b \right) \left(\int_\Omega a \right)^{\frac{2-q}{p-2}} \right)^{\frac{p-2}{p-q}} = 0,$$

which contradicts our assumption. Thus (u_n) is bounded and we may assume that $u_n \rightharpoonup u_0$. If $\lambda_s = 0$ then u_0 is a nonzero constant, and from $A(u_0) \geq 0$, $B(u_0) \geq 0$ and $S(u_0) = 1$ we get

$$\int_\Omega a \geq 0, \quad \int_{\partial\Omega} b \geq 0 \quad \text{and} \quad \int_\Omega m + \left(C_{pq}^{-1} \left(\int_{\partial\Omega} b \right) \left(\int_\Omega a \right)^{\frac{2-q}{p-2}} \right)^{\frac{p-2}{p-q}} > 0,$$

which contradicts again our assumption. Therefore $\lambda_s = \int_\Omega |\nabla u_0|^2 > 0$.

Now, let $\lambda \in (0, \lambda_s)$ and $u \in A^+ \cap B^+ \cap E_\lambda^+$. Then j_u has two critical points if $j_u(t) > 0$ for some $t > 0$. In this case, j_u has a local minimum and a global maximum, i.e. there are $t_1 < t_2$ such that $t_1 u \in \mathcal{N}_\lambda^+$ and $t_2 u \in \mathcal{N}_\lambda^-$. Note that $j_u(t) > 0$ if and only if

$i_u(t) > 0$, where i_u is given by (2.1). One may easily check that i_u has a global maximum point given by

$$t_0(u) = \left(\frac{p(2-q)}{2(p-q)} \frac{E_\lambda(u)}{\lambda A(u)} \right)^{\frac{1}{p-2}}. \tag{2.3}$$

We have

$$\begin{aligned} i_u(t_0(u)) > 0 \quad &\Leftrightarrow \quad \lambda B(u) < C_{pq} \frac{E_\lambda(u)^{\frac{p-q}{p-2}}}{(\lambda A(u))^{\frac{2-q}{p-2}}} \\ &\Leftrightarrow \quad B(u) < C_{pq} \frac{\left(\lambda^{-1}\int_\Omega |\nabla u|^2 - \int_\Omega m u^2\right)^{\frac{p-q}{p-2}}}{A(u)^{\frac{2-q}{p-2}}} \\ &\Leftrightarrow \quad E_\lambda(u) - \lambda \left(C_{pq}^{-1} B(u) A(u)^{\frac{2-q}{p-2}} \right)^{\frac{p-2}{p-q}} > 0 \\ &\Leftrightarrow \quad \int_\Omega |\nabla u|^2 - \lambda S(u) > 0, \end{aligned}$$

where $S(u)$ is defined in (1.10). Since

$$\lambda < \lambda_s = \inf \left\{ \int_\Omega |\nabla u|^2; \ u \in A_0^+ \cap B_0^+, \ S(u) = 1 \right\},$$

we get the existence of $t_1 < t_2$. Finally, from the expression of j_u it is easily seen that j_u can not have more than two critical points. □

We shall get now some kind of local coercivity for E_λ. More precisely, we shall prove that E_λ is coercive on A_0^+ (respect. B_0^+) for $\lambda < \lambda_a$ (respect. $\lambda < \lambda_b$). To this end, we deal with the maps $\alpha_1, \beta_1 : \mathbb{R} \mapsto \mathbb{R}$ given by

$$\alpha_1(\lambda) = \inf\{E_\lambda(u); \ u \in A_0^+, \ \|u\|_2 = 1\}, \tag{2.4}$$

$$\beta_1(\lambda) = \inf\{E_\lambda(u); \ u \in B_0^+, \ \|u\|_2 = 1\}. \tag{2.5}$$

Note that if $\alpha_1(\lambda) > 0$ (respect. $\beta_1(\lambda) > 0$) then $A_0^+ \setminus \{0\} \subset E_\lambda^+$ (respect $B_0^+ \setminus \{0\} \subset E_\lambda^+$). From [23], we have the following results:

Proposition 2.3.

(1) α_1 and β_1 are concave (and therefore continuous).
(2) $\lambda_b > 0$ if and only if either $\int_\Omega m < 0$ or $\int_{\partial\Omega} b < 0$. In this case, λ_b is achieved and:
  (a) $\beta_1(\lambda) > 0$ if $\lambda \in (0, \lambda_b)$. Moreover, if $\int_{\partial\Omega} b < 0$ then $\beta_1(0) > 0$.
  (b) $\lambda_b = \max\{\lambda > 0; \ \beta_1(\lambda) \geq 0\}$.
(3) Assume $\int_\Omega m < 0$. Then $\lambda_b > \lambda_1(m)$ if and only if $\int_{\partial\Omega} b\varphi_1^q < 0$.
(4) $\lambda_a > 0$ if and only if either $\int_\Omega m < 0$ or $\int_\Omega a < 0$. In this case, λ_a is achieved and:
  (a) $\alpha_1(\lambda) > 0$ if $\lambda \in (0, \lambda_a)$. Moreover, if $\int_\Omega a < 0$ then $\alpha_1(0) > 0$.
  (b) $\lambda_a = \max\{\lambda > 0; \ \alpha_1(\lambda) \geq 0\}$.
(5) Assume $\int_\Omega m < 0$. Then $\lambda_a > \lambda_1(m)$ if and only if $\int_\Omega a\varphi_1^p < 0$.

Proposition 2.4. *Assume $\int_{\partial\Omega} b < 0$. Then for every $\lambda_* \in (0, \lambda_b)$ there exist two constants $C_0 = C_0(m, b), D_0 = D_0(m, b) > 0$ such that:*

(1) $E_\lambda(u) \geq C_0 \|u\|^2$ for every $\lambda \in (0, \lambda_*)$ and $u \in B_0^+$.
(2) $B(u) \leq -D_0 \|u\|^q$ for every $\lambda \in (0, \lambda_*)$ and $u \in E_{\lambda,0}^-$.

The proof of Proposition 2.4 can be easily adapted to obtain a similar result on a:

Proposition 2.5. *Assume* $\int_\Omega a < 0$. *Then for every* $\lambda_* \in (0, \lambda_a)$ *there exist two constants* $C_0 = C_0(m,a), D_0 = D_0(m,a) > 0$ *such that:*

1. $E_\lambda(u) \geq C_0\|u\|^2$ *for every* $\lambda \in (0, \lambda_*)$ *and* $u \in A_0^+$.
2. $A(u) \leq -D_0\|u\|^p$ *for every* $\lambda \in (0, \lambda_*)$ *and* $u \in E_{\lambda,0}^-$.

Propositions 2.4 and 2.5 provide some kind of uniform coercivity (with respect to λ) to E_λ on A_0^+ and B_0^+, respectively. In case $\lambda_1(m) > 0$, we have the following:

Proposition 2.6. *Assume* $\int_\Omega m < 0$.

1. *Given any* $\mu < \lambda_1(m)$ *there exists a constant* $C_\mu > 0$ *such that* $E_\lambda(u) \geq \lambda C_\mu \|u\|^2$ *for every* $u \in H^1(\Omega)$ *and every* $\lambda \in (0, \mu)$.
2. *If* $0 < \lambda < \lambda_b$ *then there exists* $C_\lambda > 0$ *such that* $E_\lambda(u) \geq C_\lambda \|u\|^2$ *for every* $u \in B_0^+$.
3. *If* $0 < \lambda < \lambda_a$ *then there exists* $C_\lambda > 0$ *such that* $E_\lambda(u) \geq C_\lambda \|u\|^2$ *for every* $u \in A_0^+$.

Proof.

(1) Assume by contradiction that there exist $\mu < \lambda_1(m)$ and two sequences $(\lambda_n) \subset (0, \mu)$ and $(v_n) \subset H^1(\Omega)$ such that

$$E_{\lambda_n}(v_n) < \frac{\lambda_n}{n}\|v_n\|^2.$$

Setting $w_n = \frac{v_n}{\|v_n\|}$, we may assume that $w_n \rightharpoonup w_0$ in $H^1(\Omega)$ and $\lambda_n \to \lambda_* \in [0, \mu]$. Hence $E_{\lambda_*}(w_0) \leq \limsup E_{\lambda_n}(w_n) \leq 0$. If $\lambda_* > 0$ then $w_0 \equiv 0$ and $w_n \to 0$, which is impossible. If $\lambda_* = 0$ then w_0 is a constant. From

$$-\lambda_n \int_\Omega m w_n^2 < \frac{\lambda_n}{n}$$

we get $\int_\Omega m w_0^2 \geq 0$, which contradicts $\int_\Omega m < 0$.

(2) Assume that $(u_n) \subset B_0^+$ is such that

$$E_\lambda(u_n) < \frac{\|u_n\|^2}{n}.$$

Then $v_n = \frac{u_n}{\|u_n\|}$ is such that, up to a subsequence, $v_n \rightharpoonup v_0$ in $H^1(\Omega)$, $B(v_n) \to B(v_0)$, and $E_\lambda(v_n) \leq \frac{1}{n}$. Then $B(v_0) \geq 0$ and $E_\lambda(v_0) \leq 0$. Finally, $v_0 \not\equiv 0$, since otherwise we would have $v_n \to 0$, which is impossible. Therefore we get $\beta_1(\lambda) \leq 0$, which contradicts $\lambda < \lambda_b$.

(3) The proof is similar to the previous one, so we omit it.

□

3. MINIMIZATION IN $\mathcal{N}_\lambda^+$

3.1. Minimization in $\mathcal{N}_\lambda^+ \cap B^+$.

We set

$$\overline{\lambda} = \begin{cases} \min\{\lambda_s, \lambda_1\} & \text{if (2.2) holds,} \\ \min\{\lambda_s, \lambda_b\} & \text{if } \int_{\partial\Omega} b < 0 \text{ or } \int_\Omega m < 0 \text{ and } \int_\Omega a < 0. \end{cases} \tag{3.1}$$

Remark 3.1. Note that if $0 < \lambda < \overline{\lambda}$ then either $0 < \lambda < \lambda_1$ or $0 < \lambda < \lambda_b$, so that $B_0^+ \setminus \{0\} \subset E_\lambda^+$.

Let us first prove that $\mathcal{N}_\lambda^+ \cap B^+$ is non-empty and bounded for $0 < \lambda < \overline{\lambda}$:

Proposition 3.2. *Assume* $b^+ \not\equiv 0$ *and either* $\int_{\partial\Omega} b < 0$ *or* (2.2) *or* $\int_\Omega m < 0$ *and* $\int_\Omega a < 0$. *Then* $\mathcal{N}_\lambda^+ \cap B^+ \neq \emptyset$ *for every* $\lambda \in (0, \overline{\lambda})$. *Moreover:*

1. *If* (2.2) *holds then for every* $\mu < \overline{\lambda}$ *there exists a constant* $K = K_\mu > 0$ *such that* $\|u\| \leq K \|b^+\|_\infty^{\frac{1}{2-q}}$ *for every* $u \in \mathcal{N}_\lambda^+$ *and every* $0 < \lambda < \mu$.
2. *If* $\int_{\partial\Omega} b < 0$ *then for every* $\mu < \overline{\lambda}$ *there exists a constant* $K = K_\mu > 0$ *such that* $\|u\| \leq K \left(\lambda \|b^+\|_\infty\right)^{\frac{1}{2-q}}$ *for every* $u \in \mathcal{N}_\lambda^+ \cap B_0^+$ *and every* $0 < \lambda < \mu$.
3. *If* $\int_\Omega a < 0$ *and* $\int_\Omega m < 0$ *then for every* $\lambda < \overline{\lambda}$ *there exists a constant* K_λ *such that* $\|u\| \leq K_\lambda \|b^+\|_\infty^{\frac{1}{2-q}}$ *for every* $u \in \mathcal{N}_\lambda^+ \cap B_0^+$.

Proof. First of all, note that since $b^+ \not\equiv 0$ we have $B^+ \neq \emptyset$. Moreover, from Remark 3.1, if $u \in B^+$ then $u \in E_\lambda^+$. If $u \in A_0^-$ then j_u has a global minimum point $t > 0$ such that $tu \in \mathcal{N}_\lambda^+ \cap B^+$. On the other hand, if $u \in A^+$ then, by Proposition 2.2, there exists $t > 0$ such that $tu \in \mathcal{N}_\lambda^+ \cap B^+$, so that $\mathcal{N}_\lambda^+ \cap B^+ \neq \emptyset$.

1. If (2.2) holds then, given $0 < \mu < \overline{\lambda}$ and $u \in \mathcal{N}_\lambda^+$ with $0 < \lambda < \mu$, we may apply Proposition 2.6. Thus, for some $C_\mu, D > 0$, we have
$$\lambda C_\mu \|u\|^2 \leq E_\lambda(u) < \lambda \frac{p-q}{p-2} B(u) \leq \lambda D \|b^+\|_\infty \|u\|^q,$$
and consequently
$$\|u\| \leq \left(\frac{D\|b^+\|_\infty}{C_\mu} \right)^{\frac{1}{2-q}}.$$

2. If $\int_{\partial\Omega} b < 0$ then, given $0 < \mu < \overline{\lambda}$ and $u \in \mathcal{N}_\lambda^+$ with $0 < \lambda < \mu$, we apply now Proposition 2.4, so that $E_\lambda(u) \geq C_0 \|u\|^2$ for every $\lambda \in (0, \mu)$ and $u \in B_0^+$. Thus, for $u \in \mathcal{N}_\lambda^+ \cap B_0^+$ we have
$$C_0 \|u\|^2 \leq E_\lambda(u) < \lambda \frac{p-q}{p-2} B(u) \leq \lambda D \|b^+\|_\infty \|u\|^q,$$
and consequently
$$\|u\| \leq \left(\frac{\lambda D\|b^+\|_\infty}{C_0} \right)^{\frac{1}{2-q}}.$$

3. If $\int_\Omega a < 0$ and $\int_\Omega m < 0$ then, by Proposition 2.6, for every $\lambda < \lambda_b$ there exists a constant $C_\lambda > 0$ such that $E_\lambda(u) \geq C_\lambda \|u\|^2$ for $u \in B_0^+$. Thus, if $u \in \mathcal{N}_\lambda^+ \cap B_0^+$ then, for some $D > 0$ there holds
$$C_\lambda \|u\|^2 \leq E_\lambda(u) < \lambda \frac{p-q}{p-2} B(u) \leq \lambda D \|b^+\|_\infty \|u\|^q,$$
and consequently
$$\|u\| \leq \left(\frac{\lambda D\|b^+\|_\infty}{C_\lambda} \right)^{\frac{1}{2-q}}.$$

□

Remark 3.3. From the proof of Proposition 3.2 (1), we may see that if $\int_\Omega m < 0$ then for every $\mu < \lambda_1$ there exists a constant $K = K_\mu > 0$ such that $\|u\| \leq K\|b^+\|_\infty$ for every $u \in \mathcal{N}_\lambda^+$ and every $0 < \lambda < \mu$, i.e. $\mathcal{N}_\lambda^+$ is uniformly bounded for $0 < \lambda < \mu$ if $\mu < \lambda_1$.

Lemma 3.4. *Let $b, \tilde{b} \in L^\infty(\partial\Omega)$ with $b \leq \tilde{b}$. Then $\lambda_b \geq \lambda_{\tilde{b}}$ and $\lambda_s(b) \geq \lambda_s(\tilde{b})$.*

Proof. Let $B(u) = \int_{\partial\Omega} b|u|^q$ and $\tilde{B}(u) = \int_{\partial\Omega} \tilde{b}|u|^q$ for $u \in H^1(\Omega)$. Then $B(u) \leq \tilde{B}(u)$ for every $u \in H^1(\Omega)$, so that $B_0^+ \subset \tilde{B}_0^+$. It follows that $\lambda_b \geq \lambda_{\tilde{b}}$. Let us set

$$S_b(u) = \int_\Omega mu^2 + \left(C_{pq}^{-1}B(u)A(u)^{\frac{2-q}{p-2}}\right)^{\frac{p-2}{p-q}} \tag{3.2}$$

and

$$S_{\tilde{b}}(u) = \int_\Omega mu^2 + \left(C_{pq}^{-1}\tilde{B}(u)A(u)^{\frac{2-q}{p-2}}\right)^{\frac{p-2}{p-q}}. \tag{3.3}$$

Then $S_b(u) \leq S_{\tilde{b}}(u)$ for every $u \in H^1(\Omega)$. Note that we can write

$$\lambda_s(b) = \inf\left\{\frac{\int_\Omega |\nabla u|^2}{S(u)};\ u \in A_0^+ \cap B_0^+,\ S(u) > 0\right\},$$

From this formula, it follows that $\lambda_s(b) \geq \lambda_s(\tilde{b})$. □

Remark 3.5. *From Lemma 3.4 it follows that if $b_n \to b$ in $L^\infty(\partial\Omega)$ with $b^+ \not\equiv 0$ and $\int_{\partial\Omega} b < 0$ then $\overline{\lambda}(b_n)$ is bounded away from zero. Indeed, we can fix $\tilde{b} \in L^\infty(\partial\Omega)$ such that $\tilde{b}^+ \not\equiv 0$, $\int_{\partial\Omega} \tilde{b} < 0$ and $b_n \leq \tilde{b}$ for n sufficiently large. Hence $\lambda_{b_n} \geq \lambda_{\tilde{b}} > 0$ and $\lambda_s(b_n) \geq \lambda_s(\tilde{b}) > 0$ for n sufficiently large.*

Proposition 3.6. *Assume $b^+ \not\equiv 0$ and either $\int_{\partial\Omega} b < 0$ or (2.2) or $\int_\Omega m < 0$ and $\int_\Omega a < 0$. Then $\inf\limits_{\mathcal{N}_\lambda^+ \cap B^+} I_\lambda$ is achieved by some $u_{0,\lambda} \geq 0$ for $0 < \lambda < \overline{\lambda}$. Moreover:*

1. *If $\int_{\partial\Omega} b < 0$ then $u_{0,\lambda} \to 0$ in $\mathcal{C}^\theta(\overline{\Omega})$ for some $\theta \in (0,1)$ as $\lambda \to 0^+$.*
2. *If $\int_\Omega m < 0 < \int_{\partial\Omega} b$ and $\int_\Omega a < 0$ then $u_{0,\lambda} \to c_0$ in $\mathcal{C}^\theta(\overline{\Omega})$ for some $\theta \in (0,1)$ as $\lambda \to 0^+$.*
3. *If (2.2) holds then $u_{0,\lambda} \to c_1$ in $\mathcal{C}^\theta(\overline{\Omega})$ for some $\theta \in (0,1)$ as $\lambda \to 0^+$.*

Proof. Let $0 < \lambda < \overline{\lambda}$. By Proposition 3.2 we know that $\mathcal{N}_\lambda^+ \cap B^+$ is non-empty and bounded. We pick up a sequence $(u_n) \subset \mathcal{N}_\lambda^+ \cap B^+$ such that

$$I_\lambda(u_n) \to \inf_{\mathcal{N}_\lambda^+ \cap B^+} I_\lambda.$$

Since (u_n) is bounded, we may assume that

$$u_n \rightharpoonup u_0 \text{ in } H^1(\Omega) \quad \text{and} \quad B(u_n) \to B(u_0).$$

From the shape of j_u it is clear that $\inf\limits_{\mathcal{N}_\lambda^+ \cap B^+} I_\lambda < 0$. Thus $I_\lambda(u_0) < 0$, so that $u_0 \not\equiv 0$. Since

$$E_\lambda(u_n) < \lambda\frac{p-q}{p-2}B(u_n)$$

and $u_0 \in B_0^+$ we get

$$0 < E_\lambda(u_0) \leq \lambda\frac{p-q}{p-2}B(u_0),$$

i.e. $u_0 \in B^+$. From $\lambda < \lambda_s$ we deduce the existence of $t_1 > 0$ such that $t_1 u_0 \in \mathcal{N}_\lambda^+ \cap B^+$. We claim that $u_n \to u_0$. Indeed, if $u_n \not\to u_0$ then

$$j_{u_0}^{(k)}(t) < \liminf j_{u_n}^{(k)}(t)$$

for $k = 0, 1, 2$ and every $t > 0$. In particular, there holds

$$0 = j'_{u_0}(t_1) < j'_{u_n}(t_1)$$

for n sufficiently large. Thus $t_1 > 1$, since j_{u_n} is decreasing in $(0,1)$. Now, since j_{u_0} is decreasing in $(0, t_1)$ we get

$$I_\lambda(t_1 u_0) = j_{u_0}(t_1) < j_{u_0}(1) < \liminf j_{u_n}(1) = \lim I_\lambda(u_n) = \inf_{\mathcal{N}_\lambda^+ \cap B^+} I_\lambda,$$

which is a contradiction, since $t_1 u_0 \in \mathcal{N}_\lambda^+ \cap B^+$. Therefore $u_n \to u_0$ and $t_1(u_0) = 1$, so

$$I_\lambda(u_0) = \inf_{\mathcal{N}_\lambda^+ \cap B^+} I_\lambda.$$

We denote u_0 by $u_{0,\lambda}$.

(1) If $\int_{\partial\Omega} b < 0$ then, by Prop. 3.2,

$$\|u_{0,\lambda}\| \leq \left(\lambda \|b^+\|_\infty\right)^{\frac{1}{2-q}} K \to 0$$

as $\lambda \to 0^+$.

(2) If $\int_\Omega m < 0 < \int_{\partial\Omega} b$ and $\int_\Omega a < 0$, let $\lambda_n \to 0^+$ and $u_n = u_{0,\lambda_n}$. By Remark 3.3, (u_n) is bounded and we may assume that $u_n \rightharpoonup u_0$ in $H^1(\Omega)$. From

$$E_{\lambda_n}(u_n) = \lambda_n \left(A(u_n) + B(u_n)\right)$$

we infer that $\int_\Omega |\nabla u_n|^2 \to 0$, so $u_n \to u_0$ and u_0 is a constant. Since $\int_\Omega m < 0 < \int_{\partial\Omega} b$ and $\int_\Omega a < 0$, j_{u_0} has a unique critical point, which is a global minimum point. Thus there is an unique constant c in $\mathcal{N}_\lambda^+$. By Remark 2.1, we infer that $c = c_0$, where c_0 is the unique zero of φ. In particular, $c_0 \in \mathcal{N}_{\lambda_n}^+ \cap B^+$ for every n. Then

$$I_{\lambda_n}(u_n) \leq I_{\lambda_n}(c_0) < 0,$$

and consequently

$$-\frac{1}{2}\int_\Omega m u_n^2 - \frac{1}{p}A(u_n) - \frac{1}{q}B(u_n) \leq -\frac{c_0^2}{2}\int_\Omega m - \frac{c_0^p}{p}\int_\Omega a - \frac{c_0^q}{q}\int_{\partial\Omega} b < 0.$$

It follows that $u_n \not\to 0$, i.e. u_0 is a positive constant. Finally, since u_n solves (P_λ) for $\lambda = \lambda_n$, we have

$$\begin{aligned} 0 &= \int_\Omega (\nabla u_n \nabla u_0 - \lambda_n m u_n u_0) - \lambda_n \int_\Omega a u_n^{p-1} u_0 - \lambda_n \int_{\partial\Omega} b u_n^{q-1} u_0 \\ &= \lambda_n \left\{ -\int_\Omega m u_n u_0 - \int_\Omega a u_n^{p-1} u_0 - \int_{\partial\Omega} b u_n^{q-1} u_0 \right\} \end{aligned}$$

so, letting $n \to \infty$, we get

$$u_0^{2-q}\int_\Omega m + u_0^{p-q}\int_\Omega a + \int_{\partial\Omega} b = 0,$$

i.e. $u_0 = c_0$.

(3) If (2.2) holds then we can proceed as in the previous item to deduce that $u_n \to c$, where c is a constant. Now, by Remark 2.1 we infer that $c = c_1$.

□

Remark 3.7. Let $b_n \to b$ in $L^\infty(\partial\Omega)$ with $b_n^+ \not\equiv 0$ for every n, $b \not\equiv 0$ and $b \leq 0$. Then, from [23, Remark 2.5], we have $\lambda_{b_n} \to \lambda_b > 0$. Moreover, since $b_n \to b$ in $L^\infty(\partial\Omega)$, we have $\int_{\partial\Omega} b_n < 0$ for n sufficiently large and $b_n^+ \to 0$ in $L^\infty(\partial\Omega)$. So we can fix $\tilde{b} \in L^\infty(\partial\Omega)$ such that $\int_{\partial\Omega} \tilde{b} < 0$, $\tilde{b}^+ \not\equiv 0$, and $b_n \leq \tilde{b}$ for n sufficiently large. By Remark 3.5, we have $\lambda_s(b_n) \geq \lambda_s(\tilde{b}) > 0$. So $\overline{\lambda}(b_n) \geq \min\{\lambda_s(\tilde{b}), \lambda_{\tilde{b}}\}$ and u_{0,λ,b_n} exists for $0 < \lambda < \min\{\lambda_s(\tilde{b}), \lambda_{\tilde{b}}\}$ and every n. Moreover, from the proof of Proposition 3.6, for some $K_\lambda > 0$ we have $\|u_{0,\lambda,b_n}\| \leq K_\lambda \|b_n^+\|_\infty^{\frac{1}{2-q}}$, and consequently $u_{0,\lambda,b_n} \to 0$ in $\mathcal{C}^\theta(\overline{\Omega})$ for some $\theta \in (0,1)$ and $0 < \lambda < \min\{\lambda_s(\tilde{b}), \lambda_{\tilde{b}}\}$.

Lemma 3.8. *Assume $b^+ \not\equiv 0$ and $\int_{\partial\Omega} b < 0$. Then, for $0 < \lambda < \overline{\lambda}$, there holds*

$$I_\lambda(u_{0,\lambda}) < -D_0 \lambda^{\frac{2}{2-q}} + o(\lambda^{\frac{2}{2-q}}), \tag{3.4}$$

for some $D_0 > 0$.

Proof. Recall that for $0 < \lambda < \lambda_b$ we have $B^+ \subset E_\lambda^+$. Then there exists $C_0 > 0$ such that $0 < E_\lambda(u) \leq C_0 \|u\|^2$ for $u \in B^+$ and $0 < \lambda < \lambda_b$.

Let $u \in B^+ \cap A_0^+$. Then

$$I_\lambda(u) \leq \frac{1}{2} E_\lambda(u) - \frac{\lambda}{q} B(u) \leq \tilde{I}_\lambda(u) := \frac{C_0}{2}\|u\|^2 - \frac{\lambda}{q} B(u).$$

Thus $I_\lambda(tu) \leq \tilde{I}_\lambda(tu)$ for every $t > 0$. Note that $\tilde{I}_\lambda(tu)$ has a global minimum point t_0 given by

$$t_0 = \left(\frac{\lambda B(u)}{C_0 \|u\|^2} \right)^{\frac{1}{2-q}}.$$

and

$$\tilde{I}_\lambda(t_0 u) = -\frac{2-q}{2q} \lambda^{\frac{2}{2-q}} B(u) \left(\frac{B(u)}{C_0 \|u\|^2} \right)^{\frac{q}{2-q}} = -D_0 \lambda^{\frac{2}{2-q}},$$

where $D_0 = \frac{2-q}{2q} B(u) \left(\frac{B(u)}{C_0 \|u\|^2} \right)^{\frac{q}{2-q}}$. It follows that if $I_\lambda(tu)$ has a local minimum at t_1 then

$$I_\lambda(t_1 u) < -D_0 \lambda^{\frac{2}{2-q}}$$

with $D_0 > 0$

Let now $u \in B^+ \cap A^-$. Then

$$I_\lambda(u) \leq \tilde{I}_\lambda(u) := \frac{C_0}{2}\|u\|^2 - \frac{\lambda}{q} B(u) - \frac{\lambda}{p} A(u)$$

and $\tilde{I}_\lambda(tu)$ has a global minimum point t_0 which satisfies

$$t_0 C_0 \|u\|^2 - \lambda t_0^{q-1} B(u) - \lambda t_0^{p-1} A(u) = 0.$$

Thus

$$t_0 C_0 \|u\|^2 - \lambda t_0^{q-1} B(u) < 0,$$

so that

$$t_0 < \left(\frac{\lambda B(u)}{C_0 \|u\|^2} \right)^{\frac{1}{2-q}}.$$

Hence

$$\tilde{I}_\lambda(t_0 u) < -D_0 \lambda^{\frac{2}{2-q}} - \lambda^{\frac{p-q+2}{2-q}} \left(\frac{B(u)}{C_0 \|u\|^2} \right)^{\frac{1}{2-q}} A(u) = -D_0 \lambda^{\frac{2}{2-q}} + D_1 \lambda^{\frac{p-q+2}{2-q}},$$

where D_0 is as above and $D_1 = -\left(\frac{B(u)}{C_0\|u\|^2}\right)^{\frac{p}{2-q}} \frac{A(u)}{p}$. Once again, if $I_\lambda(tu)$ has a local minimum at t_1 then

$$I_\lambda(t_1 u) < -D_0\lambda^{\frac{2}{2-q}} + D_1\lambda^{\frac{p-q+2}{2-q}}.$$

Therefore we conclude that

$$\inf_{\mathcal{N}_\lambda^+ \cap B^+} I_\lambda < -D_0\lambda^{\frac{2}{2-q}} + o(\lambda^{\frac{2}{2-q}})$$

for $0 < \lambda < \overline{\lambda}$. □

Proposition 3.9. *Assume* $b^+ \not\equiv 0$ *and* $\int_{\partial\Omega} b < 0$. *If* $\lambda_n \to 0^+$ *then, up to a subsequence, there holds* $w_{0,\lambda_n} = \lambda_n^{-\frac{1}{2-q}} u_{0,\lambda_n} \to w_0$ *in* $H^1(\Omega)$, *where* w_0 *is a nontrivial non-negative solution of*

$$-\Delta w = 0 \quad \text{in } \Omega, \qquad \frac{\partial w}{\partial \mathbf{n}} = b(x)w^{q-1} \quad \text{on } \partial\Omega. \tag{3.5}$$

Proof. Let $w_n = w_{0,\lambda_n} = \lambda_n^{-\frac{1}{2-q}} u_{0,\lambda_n}$. Since

$$\|u_{0,\lambda}\| \leq \left(\lambda\|b^+\|_\infty\right)^{\frac{1}{2-q}} K$$

it follows that (w_n) is bounded in $H^1(\Omega)$. Thus, up to a subsequence, $w_n \rightharpoonup w_0$ in $H^1(\Omega)$. Furthermore, from Lemma 3.8 we have

$$I_{\lambda_n}(u_{0,\lambda_n}) < -D_0\lambda_n^{\frac{2}{2-q}} + +o(\lambda_n^{\frac{2}{2-q}}),$$

with $D_0 > 0$. Hence

$$\frac{\lambda_n^{\frac{2}{2-q}}}{2}\int_\Omega |\nabla w_n|^2 - \frac{\lambda_n^{\frac{4-q}{2-q}}}{2}\int_\Omega m w_n^2 - \frac{\lambda_n^{\frac{2}{2-q}}}{q}B(w_n) - \frac{\lambda_n^{\frac{2+p-q}{2-q}}}{p}A(w_n) < -D_0\lambda_n^{\frac{2}{2-q}} + +o(\lambda_n^{\frac{2}{2-q}}).$$

Dividing the above inequality by $\lambda_n^{\frac{2}{2-q}}$ and letting $n \to \infty$ we get

$$\frac{1}{2}\int_\Omega |\nabla w_0|^2 - \frac{1}{q}B(w_0) \leq -D_0 < 0,$$

so $w_0 \not\equiv 0$. Taking $v = w_n - w_0$ in

$$\int_\Omega \left(\nabla w_n \nabla v - \lambda_n m(x) w_n v - \lambda_n^{\frac{p-q}{2-q}} a(x) w_n^{p-1} v\right) - \int_{\partial\Omega} b(x) w_n^{q-1} v = 0 \quad \forall v \in H^1(\Omega) \tag{3.6}$$

and letting $n \to \infty$ we get $\lim \int_\Omega \nabla w_n \nabla(w_n - w_0) = 0$, so that $w_n \to w_0$ in $H^1(\Omega)$. Finally, (3.6) also shows that w_0 is a solution of (3.5). □

3.2. **Minimization in $\mathcal{N}_\lambda^+ \cap E_\lambda^-$.**

Proposition 3.10. *Assume* $\lambda_a, \lambda_b > 0$ *and* $\mathcal{N}_\lambda^+ \cap E_\lambda^- \neq \emptyset$. *If* $\inf_{\mathcal{N}_\lambda^+ \cap E_\lambda^-} I_\lambda < 0$ *and* $0 < \lambda < \min\{\lambda_a, \lambda_b\}$ *then* $\inf_{\mathcal{N}_\lambda^+ \cap E_\lambda^-} I_\lambda$ *is achieved.*

Proof. First of all, from $0 < \lambda < \min\{\lambda_a, \lambda_b\}$ we have $E_\lambda^- \subset A^- \cap B^-$ and by Proposition 2.5, taking $\lambda < \lambda_* < \min\{\lambda_a, \lambda_b\}$ we get a constant $D_0 > 0$ such that

$$A(u) \leq -D_0\|u\|^p$$

if $u \in E_\lambda^-$. Thus, from

$$\lambda A(u) = E_\lambda(u) - \lambda B(u)$$

we get

$$\lambda D_0\|u\|^p \le -\lambda A(u) < -E_\lambda(u) < \lambda C\|m\|_\infty\|u\|^2,$$

for some $C > 0$, and consequently there exists $K > 0$ such that

$$\|u\| \le K^{\frac{1}{p-2}} \tag{3.7}$$

if $u \in \mathcal{N}_\lambda^+ \cap E_\lambda^-$. Let now $(u_n) \subset \mathcal{N}_\lambda^+ \cap E_\lambda^-$ be such that

$$I_\lambda(u_n) \to \inf_{\mathcal{N}_\lambda^+ \cap E_\lambda^-} I_\lambda.$$

Since (u_n) is bounded, we may assume that $u_n \rightharpoonup u_0$ in $H^1(\Omega)$. In particular, we have $E_\lambda(u_0) \le 0$ and $I_\lambda(u_0) \le \inf_{\mathcal{N}_\lambda^+ \cap E_\lambda^-} I_\lambda < 0$, so that $u_0 \not\equiv 0$. Since $\lambda < \min\{\lambda_a, \lambda_b\}$, we have $E_{\lambda,0}^- \setminus \{0\} \subset (A^- \cap B^-) \cup \{0\}$, so $u_0 \in A^- \cap B^-$. Moreover, from

$$0 > I_\lambda(u_0) = \frac{1}{2}E_\lambda(u_0) - \frac{\lambda}{p}A(u_0) - \frac{\lambda}{q}B(u_0),$$

we infer that $E_\lambda(u_0) < 0$, i.e. $u_0 \in E_\lambda^-$. Now, as $j_{u_0}(1) = I_\lambda(u_0) < 0$, we see, from the shape of j_{u_0}, that there exists $t_2 > 0$ such that $t_2u_0 \in \mathcal{N}_\lambda^+ \cap E_\lambda^-$, i.e. t_2 is a global minimum point of j_{u_0}. If $u_n \not\to u_0$ then

$$j_{u_0}(t_2) \le j_{u_0}(1) < \liminf j_{u_n}(1) = \lim I_\lambda(u_n) = \inf_{\mathcal{N}_\lambda^+ \cap E_\lambda^-} I_\lambda,$$

which is a contradiction. Therefore $u_n \to u_0$ and consequently $t_2(u_0) = 1$. Thus $u_0 \in \mathcal{N}_\lambda^+ \cap E_\lambda^-$ and

$$I_\lambda(u_0) = \inf_{\mathcal{N}_\lambda^+ \cap E_\lambda^-} I_\lambda.$$

□

Proposition 3.11. $\mathcal{N}_\lambda^+ \cap E_\lambda^- \neq \emptyset$ *and* $\inf_{\mathcal{N}_\lambda^+ \cap E_\lambda^-} I_\lambda < 0$ *in the following cases:*

(1) $\int_\Omega m > 0 > \int_\Omega a$, $0 > \int_{\partial\Omega} b > -K_1(m,a)$ *and* $\lambda > 0$.

(2) $\int_\Omega m < 0$, $\int_{\partial\Omega} b\varphi_1^q < 0$, $-\left(\frac{C_{pq}}{-\int_{\partial\Omega} b\varphi_1^q}\right)^{\frac{p-2}{2-q}} < \int_\Omega a\varphi_1^p < 0$ *and* $\lambda > \lambda^*$, *where*

$$\lambda^* = \lambda_1(m)\left[1 - \left(C_{pq}^{-1}\left(-\int_{\partial\Omega} b\varphi_1^q\right)\left(-\int_\Omega a\varphi_1^p\right)^{\frac{2-q}{p-2}}\right)^{\frac{p-2}{p-q}}\right]^{-1}. \tag{3.8}$$

Proof. Let $u \in B^- \cap E_\lambda^- \cap A^-$. It is clear that if $j_u(t) < 0$ for some $t > 0$ then j_u has a local maximum followed by a global minimum, i.e. there are $t_1 < t_2$ such that $t_1u \in \mathcal{N}_\lambda^- \cap E_\lambda^-$ and $t_2u \in \mathcal{N}_\lambda^+ \cap E_\lambda^-$. Moreover, in this case we have $j_u(t_2) < 0$, so that $\inf_{\mathcal{N}_\lambda^+ \cap E_\lambda^-} I_\lambda < 0$. Note that $j_u(t) < 0$ if and only if $i_u(t) < 0$. Now $t_0(u)$ given by (2.3) is a global minimum point of i_u and

$$\begin{aligned} i_u(t_0(u)) < 0 \quad &\Leftrightarrow \quad \lambda(-B(u)) < C_{pq}\frac{(-E_\lambda(u))^{\frac{p-q}{p-2}}}{(-\lambda A(u))^{\frac{2-q}{p-2}}} \\ &\Leftrightarrow \quad (-B(u)) < C_{pq}\frac{\left(\int_\Omega mu^2 - \lambda^{-1}\int_\Omega |\nabla u|^2\right)^{\frac{p-q}{p-2}}}{(-A(u))^{\frac{2-q}{p-2}}} \\ &\Leftrightarrow \quad E_\lambda(u) + \lambda\left(-C_{pq}^{-1}B(u)\right)^{\frac{p-2}{p-q}}(-A(u))^{\frac{2-q}{p-q}} < 0. \end{aligned}$$

Note that if

$$\int_\Omega m > 0 > \int_\Omega a, \quad 0 > \int_{\partial\Omega} b > -K_1(m,a) \quad \text{and} \quad \lambda > 0$$

then $c \in B^- \cap E_\lambda^- \cap A^-$ for any constant c. Moreover,

$$E_\lambda(c)+\lambda\left(-C_{pq}^{-1}B(c)\right)^{\frac{p-2}{p-q}}(-A(c))^{\frac{2-q}{p-q}} = \lambda c^2\left(-\int_\Omega m + \left(-\frac{1}{C_{pq}}\int_{\partial\Omega} b\right)^{\frac{p-2}{p-q}}\left(-\int_\Omega a\right)^{\frac{2-q}{p-q}}\right),$$

so that $j_c(t_0(c)) < 0$ for $\lambda > 0$.

On the other hand, if

$$\int_\Omega m < 0, \quad \int_{\partial\Omega} b\varphi_1^q < 0, \quad -\left(\frac{C_{pq}}{-\int_{\partial\Omega} b\varphi_1^q}\right)^{\frac{p-2}{2-q}} < \int_\Omega a\varphi_1^p < 0, \quad \text{and} \quad \lambda > \lambda_1,$$

then $\varphi_1 \in B^- \cap E_\lambda^- \cap A^-$. Furthermore

$$E_\lambda(\varphi_1)+\lambda\left(-C_{pq}^{-1}B(\varphi_1)\right)^{\frac{p-2}{p-q}}(-A(\varphi_1))^{\frac{2-q}{p-q}} = \lambda_1-\lambda\left[1-\left(-\frac{1}{C_{pq}}\int_{\partial\Omega} b\varphi_1^q\right)^{\frac{p-2}{p-q}}\left(-\int_\Omega a\varphi_1^p\right)^{\frac{2-q}{p-q}}\right],$$

so that $j_{\varphi_1}(t_0(\varphi_1)) < 0$ for $\lambda > \lambda^*$.

□

Corollary 3.12. $\inf_{\mathcal{N}_\lambda^+ \cap E_\lambda^-} I_\lambda$ *is achieved by some* $u_{1,\lambda} \geq 0$ *in the following cases:*

(1) $\int_\Omega m > 0 > \int_\Omega a$, $0 > \int_{\partial\Omega} b > -K_1(m,a)$ *and* $0 < \lambda < \min\{\lambda_a, \lambda_b\}$.

(2) $\int_\Omega m < 0$, $\int_{\partial\Omega} b\varphi_1^q < 0$, $-\left(\frac{C_{pq}}{-\int_{\partial\Omega} b\varphi_1^q}\right)^{\frac{p-2}{2-q}} < \int_\Omega a\varphi_1^p < 0$ *and* $\lambda^* < \lambda < \min\{\lambda_a, \lambda_b\}$.

Remark 3.13. Let us show that the condition $\lambda^* < \min\{\lambda_a, \lambda_b\}$ assumed in Corollary 3.12 (2) may indeed hold when $\int_\Omega m < 0$. To this end, let $a_0 \in L^\infty(\Omega)$ and $b \in L^\infty(\partial\Omega)$ be such that $\int_\Omega a_0\varphi_1^p < 0$ and $\int_{\partial\Omega} b\varphi_1^q < 0$. Then $\lambda_{a_0} > \lambda_1$ and $\lambda_b > \lambda_1$. Let us set $a_\varepsilon = \varepsilon a_0$ for $\varepsilon > 0$. As one can easily see from the definition of λ_a, we have $\lambda_{a_\varepsilon} = \lambda_{a_0}$ for every $\varepsilon > 0$. Furthermore, from (3.8), we see that $\lambda^*(m, a_\varepsilon, b) \to \lambda_1$ as $\varepsilon \to 0$. Thus, for some $\varepsilon_0 > 0$ there holds $\lambda^*(m, a_\varepsilon, b) < \min\{\lambda_{a_\varepsilon}, \lambda_b\}$. The same argument applies if we consider $b_\varepsilon = \varepsilon b_0$, where $b_0 \in L^\infty(\partial\Omega)$ is such that $\int_{\partial\Omega} b_0\varphi_1^q < 0$. Therefore $\lambda^*(m, a_\varepsilon, b_\varepsilon) < \min\{\lambda_a, \lambda_b\}$ if $0 < \varepsilon < \varepsilon_0$, for some ε_0.

Proposition 3.14. *If* $\int_\Omega m > 0 > \int_\Omega a$ *and* $0 > \int_{\partial\Omega} b > -K_1(m,a)$ *then* $u_{1,\lambda} \to c_2$ *in* $\mathcal{C}^\theta(\overline{\Omega})$ *for some* $\theta \in (0,1)$ *as* $\lambda \to 0^+$.

Proof. Let $\lambda_n \to 0^+$ and $u_n = u_{1,\lambda_n}$. From (3.7) we infer that (u_n) is bounded. So, up to a subsequence, we have $u_n \rightharpoonup u_0$ in $H^1(\Omega)$. From

$$0 \geq \liminf I_{\lambda_n}(u_n) \geq \int_\Omega |\nabla u_0|^2$$

we get $u_n \to u_0$ in $H^1(\Omega)$ and u_0 is a non-negative constant. If $u_0 = 0$ then we set $v_n = \frac{u_n}{\|u_n\|}$ and assume that $v_n \rightharpoonup v_0$ in $H^1(\Omega)$. Since $v_n \in E_{\lambda_n}^-$ we have

$$\int_\Omega |\nabla v_0|^2 \leq \liminf E_{\lambda_n}(v_n) \leq 0$$

so $v_n \to v_0$ and v_0 is a positive constant. Moreover, from

$$\lambda_n\left(-\frac{1}{2}\int_\Omega mu_n^2 - \frac{1}{p}A(u_n) - \frac{1}{q}B(u_n)\right) < 0$$

we get

$$\frac{1}{q}B(v_n) \geq -\frac{1}{2}\|u_n\|^{2-q}\int_\Omega mv_n^2 - \frac{1}{p}\|u_n\|^{p-q}A(v_n).$$

Thus $B(v_0) \geq 0$, which combined with $v_n \in B^-$ provides $v_0 \in B_0$ and contradicts $\int_{\partial\Omega} b < 0$. Therefore $u_0 \neq 0$, i.e. u_0 is a positive constant. Since $u_n \to u_0$ and u_n is a solution of (P_{λ_n}) we have

$$\begin{aligned} 0 &= \int_\Omega (\nabla u_n \nabla u_0 - \lambda_n m u_n u_0) - \lambda_n \int_\Omega a u_n^{p-1} u_0 - \lambda_n \int_{\partial\Omega} b u_n^{q-1} u_0 \\ &= \lambda_n \left\{ -\int_\Omega m u_n u_0 - \int_\Omega a u_n^{p-1} u_0 - \int_{\partial\Omega} b u_n^{q-1} u_0 \right\} \end{aligned}$$

so

$$u_0^{2-q}\int_\Omega m + u_0^{p-q}\int_\Omega a + \int_{\partial\Omega} b = 0,$$

i.e. u_0 is a positive zero of φ. Finally, from $u_n \in N^+_{\lambda_n}$, we get

$$E_{\lambda_n}(u_n) < \lambda_n \frac{p-q}{p-2} B(u_n).$$

In particular, we have

$$-\lambda_n \int_\Omega m u_n^2 < \lambda_n \frac{p-q}{p-2} B(u_n),$$

so, from $u_n \to u_0$ we get

$$u_0^{2-q} \geq \frac{p-q}{p-2}\frac{\int_{\partial\Omega} b}{\left(-\int_\Omega m\right)}.$$

Since u_0 is a positive zero of φ, we get

$$u_0 \geq \left(-\frac{2-q}{p-q}\frac{\int_\Omega m}{\int_\Omega a}\right)^{\frac{1}{p-2}},$$

so that, by (1.6), $u_0 = c_2$. ☐

4. Minimization in $\mathcal{N}_\lambda^-$

Proposition 4.1. *Assume* $a^+ \not\equiv 0$ *and either* $\int_\Omega a < 0$ *or* (2.2). *Then* $\mathcal{N}_\lambda^- \cap A^+ \neq \emptyset$ *for every* $0 < \lambda < \min\{\lambda_a, \lambda_s\}$. *Moreover, for* $0 < \lambda < \min\{\lambda_a, \lambda_s\}$, *there holds:*

1. $\mathcal{N}_\lambda^- \cap A^+$ *is bounded away from zero, i.e. there exists* $K_\lambda > 0$ *such that* $\|u\| \geq K_\lambda$ *for* $u \in \mathcal{N}_\lambda^- \cap A^+$.
2. *If* $(u_n) \subset \mathcal{N}_\lambda^- \cap A^+$ *is a sequence such that* $(I_\lambda(u_n))$ *is bounded from above then* (u_n) *is bounded.*
3. $\inf_{\mathcal{N}_\lambda^- \cap A^+} I_\lambda > 0$.

Proof. First of all, note that since $a^+ \not\equiv 0$ we have $A^+ \neq \emptyset$. Let $u \in A^+$. If $u \in B^-$ then j_u has a global maximum point $t_1 > 0$, so $t_1 u \in \mathcal{N}_\lambda^-$. The same conclusion holds if $u \in B_0^+$, since $\lambda < \lambda_a$ provides $u \in E_\lambda^+$, whereas $0 < \lambda < \lambda_s$ yields that j_u has a global maximum point, by Proposition 2.2. Therefore $\mathcal{N}_\lambda^- \cap A^+ \neq \emptyset$.

(1) Let us assume first $\int_\Omega a < 0$. Given $0 < \lambda < \mu < \lambda_a$ and $u \in \mathcal{N}_\lambda^- \cap A^+$, we apply Proposition 2.5. Then, for some $C_0, D > 0$, we have

$$C_0\|u\|^2 \leq E_\lambda(u) < \lambda\frac{p-q}{2-q}A(u) \leq \lambda D\|a^+\|_\infty \|u\|^p, \tag{4.1}$$

and consequently

$$\|u\| \geq \left(\frac{C_0}{\lambda D\|a^+\|_\infty}\right)^{\frac{1}{p-2}}. \tag{4.2}$$

Now, if (2.2) holds then, since $\lambda < \lambda_a$, there is a constant $C_\lambda > 0$ such that $E_\lambda(u) \geq C_\lambda\|u\|^2$ for every $u \in A^+$. Thus

$$C_\lambda\|u\|^2 \leq E_\lambda(u) < \lambda\frac{p-q}{2-q}A(u) \leq \lambda D\|a^+\|_\infty\|u\|^p,$$

and consequently

$$\|u\| \geq \left(\frac{C_\lambda}{\lambda D\|a^+\|_\infty}\right)^{\frac{1}{p-2}}$$

if $u \in \mathcal{N}_\lambda^- \cap A^+$.

(2) Note that

$$I_\lambda(u) = \frac{p-2}{2p}E_\lambda(u) - \lambda\frac{p-q}{pq}B(u)$$

if $u \in \mathcal{N}_\lambda^-$. Hence, as $\lambda < \lambda_a$, if in addition $u \in A^+$ then

$$I_\lambda(u) \geq C_\lambda\|u\|^2 - \lambda D\|u\|^q,$$

for some constants $C_\lambda, D > 0$. From the above inequality we deduce that if $(u_n) \subset \mathcal{N}_\lambda^- \cap A^+$ is such that $(I_\lambda(u_n))$ is bounded from above then (u_n) is bounded.

(3) If $u \in \mathcal{N}_\lambda^-$ then

$$I_\lambda(u) = \frac{p-2}{2p}E_\lambda(u) - \lambda\frac{p-q}{qp}B(u). \tag{4.3}$$

If, in addition, $u \in A^+$ then $E_\lambda(u) \geq C_\lambda\|u\|^2$ for some constant $C_\lambda > 0$, since $\lambda < \lambda_a$. Thus, if $u \in \mathcal{N}_\lambda^- \cap A^+ \cap B_0^-$ then

$$I_\lambda(u) \geq D_\lambda\|u\|^2 \geq \tilde{D}_\lambda > 0,$$

where we used (1). Now, if $(u_n) \subset \mathcal{N}_\lambda^- \cap A^+ \cap B^+$ then, from $\lambda < \lambda_s$, we have

$$I_\lambda(u_n) = j_{u_n}(1) \geq j_{u_n}(t_0(u_n)) > 0.$$

If, in addition $I_\lambda(u_n) \to 0$ then $j_{u_n}(t_0(u_n)) \to 0$, and consequently either $t_0(u_n) \to 0$ or $i_{u_n}(t_0(u_n)) \to 0$. In the first case, we get $E_\lambda(u_n) \to 0$, so that $u_n \to 0$ in $H^1(\Omega)$, which contradicts (1). Now, if $i_{u_n}(t_0(u_n)) \to 0$ then

$$-\lambda B(u_n) + C_{pq}\frac{E_\lambda(u_n)^{\frac{p-q}{p-2}}}{(\lambda A(u_n))^{\frac{2-q}{p-2}}} \to 0^+.$$

Since (u_n) is bounded, we may assume that

$$u_n \rightharpoonup u_0 \text{ in } H^1(\Omega), \quad A(u_0) = \lim A(u_n) \geq 0 \quad \text{and} \quad B(u_0) = \lim B(u_n).$$

One may easily see that if either $B(u_0) = 0$ or $A(u_0) = 0$ then $E_\lambda(u_n) \to 0$ and we infer again that $u_n \to 0$ in $H^1(\Omega)$, which contradicts (1). Thus $u_0 \in A^+ \cap B^+ \cap E_\lambda^+$. From $\lambda < \lambda_s$ we have

$$0 < i_{u_0}(t_0(u_0)) \leq \lim i_{u_n}(t_0(u_n)),$$

which is a contradiction. Therefore we can't have $I_\lambda(u_n) \to 0$, so $\inf_{\mathcal{N}_\lambda^- \cap A^+} I_\lambda > 0$.

□

Remark 4.2.

(1) If (2.2) holds and $\lambda < \lambda_1$ then the conclusions of Proposition 4.1 remain valid. Indeed, in this case, for every $\lambda < \lambda_1$ there exists a constant $C_\lambda > 0$ such that $E_\lambda(u) \geq C_\lambda \|u\|^2$ for every $u \in H^1(\Omega)$.
(2) If $\int_\Omega a < 0$ then (4.2) shows that $\mathcal{N}_\lambda^- \cap A^+$ is uniformly bounded away from zero for $\lambda \in (0, \mu)$, with $\mu < \lambda_a$. Furthermore, the statement in (2) can be strengthened as follows: if $(\lambda_n) \subset (0, \mu)$ and $(u_n) \subset \mathcal{N}_{\lambda_n}^- \cap A^+$ are such that $I_{\lambda_n}(u_n)$ is bounded then (u_n) is bounded. As a matter of fact, in this case we have
$$I_{\lambda_n}(u_n) \geq C_0 \|u_n\|^2 - \mu D \|u_n\|^q$$
for some constants $C_0, D > 0$.

Proposition 4.3. *Assume* $a^+ \not\equiv 0$. *If either* (2.2) *or* $\int_\Omega a < 0$ *holds then* $\inf_{\mathcal{N}_\lambda^- \cap A^+} I_\lambda$ *is achieved by some* $u_{2,\lambda} \geq 0$ *for* $0 < \lambda < \min\{\lambda_s, \lambda_a\}$. *Moreover, if* (2.2) *holds then* $u_{2,\lambda} \to c_2$ *in* $\mathcal{C}^\theta(\overline{\Omega})$ *for some* $\theta \in (0,1)$ *as* $\lambda \to 0^+$.

Proof. Let $0 < \lambda < \min\{\lambda_s, \lambda_a\}$ and $(u_n) \subset \mathcal{N}_\lambda^- \cap A^+$ be such that
$$I_\lambda(u_n) \to \inf_{\mathcal{N}_\lambda^- \cap A^+} I_\lambda.$$
By Proposition 4.1, we know that (u_n) is bounded, so we may assume that $u_n \rightharpoonup u_0$ in $H^1(\Omega)$ and $u_0 \in A_0^+$. From
$$\lambda \frac{p-q}{p-2} B(u_n) < E_\lambda(u_n) < \lambda \frac{p-q}{2-q} A(u_n) \tag{4.4}$$
we have
$$E_\lambda(u_0) \leq \liminf E_\lambda(u_n) \leq \limsup E_\lambda(u_n) \leq \lambda \frac{p-q}{2-q} A(u_0).$$
If $u_0 \equiv 0$ then the above inequalities provide $E_\lambda(u_n) \to E_\lambda(u_0)$, so $u_n \to u_0 \equiv 0$. This is impossible by Proposition 4.1 (1). Hence $u_0 \not\equiv 0$. Moreover, if $A(u_0) = 0$ then $E_\lambda(u_0) \leq 0$, with $u_0 \not\equiv 0$, which contradicts $\lambda < \lambda_a$. Thus $u_0 \in A^+$ and it is easily seen that there exists $t_2 > 0$ such that $t_2 u_0 \in \mathcal{N}_\lambda^- \cap A^+$. We claim that $u_n \to u_0$. Indeed, if not then
$$j_{u_0}(t) < \liminf j_{u_n}(t)$$
for every $t > 0$. Hence
$$I_\lambda(t_2 u_0) = j_{u_0}(t_2) < \liminf j_{u_n}(t_2) \leq \liminf j_{u_n}(1) = \lim I_\lambda(u_n) = \inf_{\mathcal{N}_\lambda^- \cap A^+} I_\lambda.$$
We have then a contradiction, so $u_n \to u_0$ in $H^1(\Omega)$ and
$$I_\lambda(u_0) = \inf_{\mathcal{N}_\lambda^- \cap A^+} I_\lambda.$$
We denote u_0 by $u_{2,\lambda}$.

Assume now (2.2). Let $\lambda_n \to 0^+$ and $u_n = u_{2,\lambda_n}$. We claim that (u_n) is bounded. Indeed, assume that $\|u_n\| \to \infty$ and set $v_n = \frac{u_n}{\|u_n\|}$. We may assume that $v_n \rightharpoonup v_0$ in $H^1(\Omega)$. Note that since $\int_\Omega a > 0$ and $0 < \lambda < \lambda_s$, there is a unique positive constant $c \in \mathcal{N}_{\lambda_n}^- \cap A^+$ for every n. Thus
$$\frac{p-2}{2p} \lambda_n A(u_n) - \frac{2-q}{2q} \lambda_n B(u_n) = I_{\lambda_n}(u_n) \leq \lambda_n \left(-\frac{\mathcal{C}^2}{2} \int_\Omega m - \frac{\mathcal{C}^p}{p} \int_\Omega a - \frac{\mathcal{C}^q}{q} \int_{\partial\Omega} b \right).$$
It follows that $I_{\lambda_n}(u_n) \to 0$ and $A(v_0) = \lim A(v_n) = 0$. If $u_n \in B_0^-$ then
$$I_{\lambda_n}(u_n) \geq \frac{p-2}{2p} E_{\lambda_n}(u_n),$$

so that $E_{\lambda_n}(v_n) \to 0$, and consequently $\int_\Omega |\nabla v_n|^2 \to 0$, i.e. v_0 is a nonzero constant. This contradicts $A(v_0) = 0$. On the other hand, if $u_n \in B^+$ then $u_n \in B^+ \cap A^+$ and $j_{u_n}(t_0(u_n)) \to 0$. So either $t_0(u_n) \to 0$ or

$$-\lambda_n B(v_n) + C_{pq} \frac{E_{\lambda_n}(v_n)^{\frac{p-q}{p-2}}}{(\lambda_n A(v_n))^{\frac{2-q}{p-2}}} \to 0^+.$$

Since

$$\begin{aligned} t_0(u_n) &= \left(\frac{p(2-q)}{2(p-q)} \frac{E_{\lambda_n}(u_n)}{\lambda_n A(u_n)}\right)^{\frac{1}{p-2}} = \left(\frac{p(2-q)}{2(p-q)} \left(1 + \frac{B(u_n)}{A(u_n)}\right)\right)^{\frac{1}{p-2}} \\ &> \left(\frac{p(2-q)}{2(p-q)}\right)^{\frac{1}{p-2}}, \end{aligned}$$

the first case is ruled out. In the second case, it follows that $E_{\lambda_n}(v_n) \to 0$ and once again we deduce that v_0 is a nonzero constant, which is impossible. Therefore (u_n) is bounded and we may assume that $u_n \rightharpoonup u_0$ in $H^1(\Omega)$. From

$$E_{\lambda_n}(u_n) = \lambda_n \left(A(u_n) + B(u_n)\right)$$

we infer that $\int_\Omega |\nabla u_n|^2 \to 0$, so $u_n \to u_0$ and u_0 is a constant. Since $\|u_n\| \geq K > 0$, we know that $u_0 \neq 0$. Finally, proceeding as in the proof of Proposition 3.14, we see that $u_0 = c_2$. $\square$

Remark 4.4. Let $a_n \to a$ in $L^\infty(\Omega)$ with $a_n^+ \not\equiv 0$ for every n, $a \not\equiv 0$ and $a \leq 0$. Arguing as in the proof of Lemma 3.4, we may show that if $\tilde{a} \in L^\infty(\Omega)$ is such that $\int_\Omega \tilde{a} < 0$ and $\tilde{a} \geq a_n$ for every n, then $\lambda_{a_n} \geq \lambda_{\tilde{a}}$ and $\lambda_s(a_n) \geq \lambda_s(\tilde{a})$ for every n. So u_{2,λ,a_n} exists for $0 < \lambda < \min\{\lambda_s(\tilde{a}), \lambda_{\tilde{a}}\}$ and every n. Moreover, since $a_n^+ \to 0$ in $L^\infty(\Omega)$, from (4.2) we have $\|u_{2,\lambda,a_n}\| \to \infty$ for $0 < \lambda < \min\{\lambda_s(\tilde{a}), \lambda_{\tilde{a}}\}$. Finally, getting back to (4.1), we deduce that $\int_\Omega a_n u_{2,\lambda,a_n}^p \to \infty$ and therefore $\|u_{2,\lambda,a_n}\|_{\mathcal{C}(\overline{\Omega})} \to \infty$ for $0 < \lambda < \min\{\lambda_s(\tilde{a}), \lambda_{\tilde{a}}\}$.

Proposition 4.5. *Assume $a^+ \not\equiv 0$ and $\int_\Omega a < 0$. If $\lambda_n \to 0^+$ then, up to a subsequence, there holds $w_n := \lambda_n^{\frac{1}{p-2}} u_{2,\lambda_n} \to w_\infty$ in $H^1(\Omega)$, where w_∞ is a nontrivial non-negative solution of the problem*

$$-\Delta w = a(x) w^{p-1} \quad \text{in } \Omega, \qquad \frac{\partial w}{\partial \mathbf{n}} = 0 \quad \text{on } \partial\Omega. \tag{4.5}$$

Proof. We claim that (w_n) is bounded in $H_0^1(\Omega)$. Indeed, note that w_n minimizes J_{λ_n} over $\mathcal{M}_{\lambda_n}^- \cap A^+$, where

$$J_\lambda(w) = \frac{1}{2} E_\lambda(w) - \frac{1}{q} \lambda^{\frac{p-q}{p-2}} B(w) - \frac{1}{p} A(w) \quad \text{for } w \in H^1(\Omega),$$

and $\mathcal{M}_\lambda$ is the Nehari manifold associated to J_λ. If $w \in \mathcal{M}_\lambda^- \cap A^+$ and $0 < \lambda < \min\{\lambda_a, \lambda_s\}$ then

$$J_\lambda(w) = \left(\frac{1}{2} - \frac{1}{p}\right) E_\lambda(w) - \left(\frac{1}{q} - \frac{1}{p}\right) \lambda^{\frac{p-q}{p-2}} B(w) \geq C_0 \|w\|^2 - C_1 \|w\|^q$$

for some constants $C_0, C_1 > 0$. If we prove that $J_{\lambda_n}(w_n)$ is bounded from above then we deduce that (w_n) is bounded. We have

$$J_{\lambda_n}(w_n) = \inf_{\mathcal{M}_{\lambda_n}^- \cap A^+} J_{\lambda_n} \leq \inf_{\mathcal{M}_{\lambda_n}^- \cap A^+ \cap H_0^1(\Omega)} J_{\lambda_n}.$$

If $w \in H_0^1(\Omega)$ then $B(w) = 0$, so $J_\lambda(w) = \frac{1}{2} E_\lambda(w) - \frac{1}{p} A(w)$, and it can be shown that $\inf_{\mathcal{M}_\lambda^- \cap A^+ \cap H_0^1(\Omega)} J_\lambda$ is achieved for $\lambda \in (0, \lambda_1^D(m))$, where $\lambda_1^D(m)$ is the first positive eigenvalue of

$$-\Delta u = \lambda m(x)u \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega.$$

Finally, we claim that the latter infimum is bounded from above for $\lambda \in (0, \lambda_1^D(m))$, which yields the conclusion. This claim follows from the inequality

$$J_\lambda(w) \le L(w) := \frac{C}{2}\|w\|^2 - \frac{1}{p}A(w),$$

which holds for $w \in H_0^1(\Omega)$, $\lambda \in (0, \lambda_1^D(m))$ and some $C > 0$. Thus, given $w \in H_0^1(\Omega) \cap A^+$, if $J_\lambda(tw)$ achieves its global maximum at $t_0 > 0$ then

$$J_\lambda(t_0 w) \le L(t_0 w) \le \left(\frac{1}{2} - \frac{1}{p}\right)\left(\frac{C\|w\|^2}{A(w)}\right)^{\frac{1}{p-2}} C\|w\|^2.$$

Therefore, fixing a $w_0 \in H_0^1(\Omega) \cap A^+$, we obtain

$$\inf_{\mathcal{M}_\lambda^- \cap A^+ \cap H_0^1(\Omega)} J_\lambda \le K := \left(\frac{1}{2} - \frac{1}{p}\right)\left(\frac{C\|w_0\|^2}{A(w_0)}\right)^{\frac{1}{p-2}} C\|w_0\|^2,$$

for $\lambda \in (0, \lambda_1^D(m))$, as claimed. Thus (w_n) is bounded in $H^1(\Omega)$, so up to a subsequence we have $w_n \rightharpoonup w_\infty$ in $H^1(\Omega)$. Taking $v = w_n - w_\infty$ in

$$\int_\Omega \left(\nabla w_n \nabla v - \lambda_n m(x) w_n v - a(x) w_n^{p-1} v\right) - \lambda_n^{\frac{p-q}{p-2}} \int_{\partial\Omega} b(x) w_n^{q-1} v = 0 \quad \forall v \in H^1(\Omega) \tag{4.6}$$

and letting $n \to \infty$ we get $\lim \int_\Omega \nabla w_n \nabla (w_n - w_\infty) = 0$, so that $w_n \to w_\infty$ in $H^1(\Omega)$. Furthermore, since

$$C_0\|w_n\|^2 \le E_{\lambda_n}(w_n) < \frac{p-q}{2-q} A(w_n) \le C_1 \|w_n\|^p$$

for some $C_0, C_1 > 0$, we get $\|w_n\| \ge C^{\frac{1}{p-2}}$ for some $C > 0$, so that $w_\infty \not\equiv 0$. Finally, (4.6) also shows that w_∞ is a solution of (4.5). ☐

5. Proofs of the main results

Before proceeding to the proofs of our main results, we prove a partial positivity result on the boundary for nontrivial non-negative solutions of (P_λ):

Proposition 5.1.

(1) *Let* u_λ *be a nontrivial non-negative solution of* (P_λ) *for* $\lambda > 0$. *Then the set* $\{x \in \partial\Omega : u(x) = 0\}$ *has no interior points in the relative topology of* $\partial\Omega$, *and it is contained in* $\{x \in \partial\Omega : b(x) \le 0\}$ *if* $b \in \mathcal{C}(\partial\Omega)$.

(2) *Let* w_0 *be a nontrivial non-negative solution of* (1.12). *Then* $w_0 > 0$ *in* Ω, *the set* $\{x \in \partial\Omega : w_0 = 0\}$ *has no interior points in the relative topology of* $\partial\Omega$, *and it is contained in* $\{x \in \partial\Omega : b(x) \le 0\}$ *if* $b \in \mathcal{C}(\partial\Omega)$.

Proof.

(1) Assume by contradiction that x_0 is an interior point of $\partial\Omega$ with $u_\lambda(x_0) = 0$. Then, there exists $\rho_0 > 0$ such that $u_\lambda(x) = 0$ for $x \in \Gamma_1 := B_{\rho_0}(x_0) \cap \partial\Omega$. Let D be a

subdomain of Ω with smooth boundary ∂D such that $\Gamma_1 \subset \partial D$ and $\Gamma_0 := \partial D \setminus \overline{\Gamma_1} = \partial D \cap \Omega$. Consider the following mixed problem

$$\begin{cases} -\Delta u = \lambda(-m_\infty u - a_\infty u^{p-1}) & \text{in } D, \\ \frac{\partial u}{\partial \mathbf{n}} = \lambda K_\infty u & \text{on } \Gamma_1, \\ u = 0 & \text{on } \Gamma_0, \end{cases} \tag{5.1}$$

where $m_\infty = \|m\|_\infty$, $a_\infty = \|a\|_\infty > 0$, and $K_\infty > 0$ is a constant to be determined. Arguing as in the proof of [16, Theorem 1], we can prove that if K_∞ is sufficiently large then (5.1) has a unique nontrivial non-negative weak solution $v_\lambda \in H^1_{\Gamma_0}(D)$. Here, $H^1_{\Gamma_0}(D)$ is defined as the closure of $\mathcal{C}^\infty_c\left(D \cup (\partial D \setminus \overline{\Gamma_0})\right)$ with respect to the $H^1(D)$ norm. We remark that $v_\lambda \in \mathcal{C}^2(D \cup \Gamma_1) \cap \mathcal{C}(\overline{D})$ [1, 26], so that $v_\lambda > 0$ in $D \cup \Gamma_1$ by the strong maximum principle and the boundary point lemma. On the other hand, we have $u_\lambda > 0$ on Γ_0, and for any $\varphi \in H^1_{\Gamma_0}(D)$ satisfying $\varphi \geq 0$ there holds

$$\int_D \nabla u_\lambda \nabla \varphi - \lambda \int_D \left(-m_\infty u_\lambda - a_\infty u_\lambda^{p-1}\right) \varphi - \lambda \int_{\Gamma_1} K_\infty u_\lambda \varphi \geq 0,$$

since $u_\lambda = 0$ on Γ_1. Hence, by Proposition A.1, we deduce that $v_\lambda \leq u_\lambda$ in $\overline{D}$. Thus $u_\lambda(x_0) = 0 < v_\lambda(x_0)$, and a contradiction follows.

The second assertion can be verified in a similar way. We assume that $u_\lambda(x_0) = 0$ but $b(x_0) > 0$ for some $x_0 \in \partial\Omega$. Then there exist $\rho_0, b_0 > 0$ such that $b(x) \geq b_0$ for $x \in \Gamma_1 := B_{\rho_0}(x_0) \cap \partial\Omega$. Setting D, Γ_0 as above, we consider the following mixed problem

$$\begin{cases} -\Delta u = \lambda(-m_\infty u - a_\infty u^{p-1}) & \text{in } D, \\ \frac{\partial u}{\partial \mathbf{n}} = \lambda b_0 u^{q-1} & \text{on } \Gamma_1, \\ u = 0 & \text{on } \Gamma_0. \end{cases} \tag{5.2}$$

By direct computations, we have

$$\int_D \nabla u_\lambda \nabla \varphi - \lambda \int_D \left(-m_\infty u_\lambda - a_\infty u_\lambda^{p-1}\right) \varphi - \lambda \int_{\Gamma_1} b_0 u_\lambda^{q-1} \varphi \geq 0$$

for any $\varphi \in H^1_{\Gamma_0}(D)$ satisfying $\varphi \geq 0$. Moreover, we have $u_\lambda > 0$ in $\Gamma_0 \cup D$, and $u_\lambda \in \mathcal{C}^\theta(\overline{D})$ for some $\theta \in (0,1)$. On the other hand, associated with (5.2), we consider the following eigenvalue problem.

$$\begin{cases} -\Delta \phi = \lambda(-m_\infty)\phi + \sigma\phi & \text{in } D, \\ \frac{\partial \phi}{\partial \mathbf{n}} = \lambda K \phi & \text{on } \Gamma_1, \\ \phi = 0 & \text{on } \Gamma_0. \end{cases} \tag{5.3}$$

We note that if $K > 0$ is sufficiently large then for every $\lambda > 0$ the above problem has a negative first eigenvalue σ_1, cf. [16]. Let ϕ_1 be the positive eigenfunction associated to σ_1 with $\|\phi_1\|_\infty = 1$. Since $\phi_1 \in \mathcal{C}^2(\Omega \cup \Gamma_1) \cap \mathcal{C}(\overline{D})$, by the strong maximum principle and the boundary point lemma, we have $\phi_1 > 0$ in $D \cup \Gamma_1$. By direct computations, if $\varepsilon > 0$ is a constant then, for any $\varphi \in H^1_{\Gamma_0}(D)$ satisfying $\varphi \geq 0$, we have

$$\int_D \nabla(\varepsilon\phi_1)\nabla\varphi - \lambda \int_D \left(-m_\infty \varepsilon\phi_1 - a_\infty (\varepsilon\phi_1)^{p-1}\right) \varphi - \lambda \int_{\Gamma_1} b_0 (\varepsilon\phi_1)^{q-1} \varphi \leq 0,$$

provided that

$$0 < \varepsilon \leq \min\left\{ \left(\frac{-\sigma_1}{\lambda a_\infty}\right)^{\frac{1}{p-2}}, \left(\frac{b_0}{K}\right)^{\frac{1}{2-q}} \right\}.$$

Applying Proposition A.1 to (5.2) with $u = \varepsilon\phi_1$ and $v = u_\lambda$, we obtain $\varepsilon\phi_1 \leq u_\lambda$ in $\overline{D}$. However, we have $u_\lambda(x_0) = 0 < \varepsilon\phi_1(x_0)$, which is a contradiction.

(2) First of all, by the weak maximum principle, we have $w_0 > 0$ in Ω. We argue now as in the previous item to deduce the positivity result on $\partial\Omega$. As a matter of fact, it suffices to consider (5.1) replaced by the problem

$$\begin{cases} -\Delta u = \lambda(w_0 u - u^2) & \text{in } D, \\ \frac{\partial u}{\partial \mathbf{n}} = 0 & \text{on } \Gamma_1, \\ u = 0 & \text{on } \Gamma_0, \end{cases} \tag{5.4}$$

and note that (5.4) has a unique nontrivial non-negative weak solution for $\lambda > 0$ large. Note also that if b is continuous and $b(x_0) > 0 = w(x_0)$ then we can apply the same argument to reach a contradiction, so that $\{x \in \partial\Omega : w_0 = 0\} \subset \{x \in \partial\Omega : b(x) \leq 0\}$.

□

Remark 5.2. *If u_λ is a nontrivial non-negative solution of (P_λ) for $\lambda < 0$ then the assertions of Proposition 5.1 (1) hold true replacing $\{x \in \partial\Omega : b(x) \leq 0\}$ by $\{x \in \partial\Omega : b(x) \geq 0\}$. Indeed, if $\lambda < 0$ then, by the change of variables $\mu = -\lambda$, (P_λ) reduces to*

$$\begin{cases} -\Delta u = \mu\left((-m)u + (-a)|u|^{p-2}u\right) & \textit{in } \Omega, \\ \frac{\partial u}{\partial \mathbf{n}} = \mu(-b)|u|^{q-2}u & \textit{on } \partial\Omega. \end{cases}$$

with $\mu > 0$.

We deduce now our existence results using the fact that local minimizers of I_λ constrained to $\mathcal{N} \setminus \mathcal{N}_0$ are critical points of I_λ and, therefore, solutions of (P_λ). It is clear that A^+, B^+ and E_λ^- are open sets, so that, whenever achieved, the infima of I_λ constrained to $\mathcal{N}_\lambda^+ \cap B^+$, $\mathcal{N}_\lambda^+ \cap E_\lambda^-$ and $\mathcal{N}_\lambda^- \cap A^+$ provide solutions of (P_λ).

Proof of Theorem 1.3:

By Propositions 2.2 and 2.3, we have $\lambda_b, \lambda_s > 0$ if $\int_{\partial\Omega} b < 0$ and $\lambda_a, \lambda_s > 0$ if $\int_\Omega a < 0$. Moreover, if $\int_{\partial\Omega} b < 0$ then Proposition 3.6 yields that $\inf\limits_{\mathcal{N}_\lambda^+ \cap B^+} I_\lambda$ is achieved by $u_{0,\lambda} \geq 0$ for $0 < \lambda < \min\{\lambda_b, \lambda_s\}$ and $u_{0,\lambda} \to 0$ in $\mathcal{C}^\theta(\overline{\Omega})$ for some $\theta \in (0,1)$ as $\lambda \to 0^+$. Likewise, if $\int_\Omega a < 0$ then, by Proposition 4.3, $\inf\limits_{\mathcal{N}_\lambda^- \cap A^+} I_\lambda$ is achieved by $u_{2,\lambda} \geq 0$ for $0 < \lambda < \min\{\lambda_a, \lambda_s\}$. Furthermore, by Propositions 3.9 and 4.5, the asymptotic profiles of $u_{0,\lambda}$ and $u_{2,\lambda}$ are given by $\lambda^{\frac{1}{q-2}} w_0$ and $\lambda^{\frac{1}{p-2}} w_\infty$ as $\lambda \to 0^+$, where w_0 and w_∞ are nontrivial non-negative solutions of (3.5) and (4.5), respectively.

By a standard bootstrap argument, we obtain $w_\infty \in W^{2,r}(\Omega)$, with $r > N$. The strong maximum principle and boundary point lemma yield $w_\infty > 0$ in $\overline{\Omega}$. Setting $w_\lambda = \lambda^{\frac{1}{p-2}} u_{2,\lambda}$, we have that w_λ is bounded in $H^1(\Omega)$ and w_λ is a weak solution of the problem

$$\begin{cases} -\Delta w = \lambda m w + a w^{p-1} & \text{in } \Omega, \\ \frac{\partial w}{\partial \mathbf{n}} = \lambda^{\frac{p-q}{p-2}} b w^{q-1} & \text{on } \partial\Omega. \end{cases}$$

Rossi's bootstrap argument [25] yields that w_λ is bounded in $\mathcal{C}^\nu(\overline{\Omega})$ for some $\nu \in (0,1)$. By the compact embedding $\mathcal{C}^\nu(\overline{\Omega}) \subset \mathcal{C}^\theta(\overline{\Omega})$, $\theta < \nu$, we may obtain that w_λ converges to some

w^* in $\mathcal{C}^{\theta}(\overline{\Omega})$. Since $w_\lambda \to w_\infty$ in $L^2(\Omega)$, we have $w^* = w_\infty$. From $w_\infty > 0$ in $\overline{\Omega}$, we have $u_{2,\lambda} > 0$ in $\overline{\Omega}$ for $\lambda > 0$ close to 0, and $\min_{\overline{\Omega}} u_{2,\lambda} \to \infty$ as $\lambda \to 0^+$. It can be verified in a similar way that $\lambda^{-\frac{1}{2-q}} u_{0,\lambda} \to w_0$ in $C^{\tilde{\theta}}(\overline{\Omega})$ for some $\tilde{\theta} \in (0,1)$. Indeed, it suffices to note that $\lambda^{-\frac{1}{2-q}} u_{0,\lambda}$ is a weak solution of the problem

$$\begin{cases} -\Delta w = \lambda m w + \lambda^{\frac{p-q}{2-q}} a w^{p-1} & \text{in } \Omega, \\ \frac{\partial w}{\partial \mathbf{n}} = b w^{q-1} & \text{on } \partial\Omega. \end{cases}$$

Finally, Proposition 5.1 provides the positivity properties of w_0. $\square$

Proof of Theorem 1.5:

By Corollary 3.12, $\inf\limits_{\mathcal{N}_\lambda^+ \cap E_\lambda^-} I_\lambda$ is achieved by $u_{1,\lambda} \geq 0$ for $0 < \lambda < \min\{\lambda_a, \lambda_b\}$. Moreover, by Proposition 3.14, $u_{1,\lambda} \to c_2$ in $\mathcal{C}^{\theta}(\overline{\Omega})$ for some $\theta \in (0,1)$ as $\lambda \to 0^+$.

Now, if $\lambda < 0$ then we change the signs of λ, m, a and b. Since

$$\int_\Omega (-m) < 0 < \int_\Omega (-a) \quad \text{and} \quad 0 < \int_{\partial\Omega} (-b) < K_1(m,a),$$

Propositions 3.6 and 4.3 yield the existence of two non-negative solutions $u_{0,-\lambda}$, $u_{2,-\lambda}$ for $0 < -\lambda < \min\{\tilde{\lambda}_1, \tilde{\lambda}_s\}$, which satisfy $u_{0,-\lambda} \to c_1$ and $u_{2,-\lambda} \to c_2$ in $\mathcal{C}^{\theta}(\overline{\Omega})$ for some $\theta \in (0,1)$ as $\lambda \to 0^-$. We set then $v_{1,\lambda} = u_{0,-\lambda}$ and $v_{2,\lambda} = u_{2,-\lambda}$. $\square$

Proof of Theorem 1.6:

First of all, by a standard bootstrap argument we infer that $u_{2,\lambda}$ is a classical positive solution of (P_λ), since $u_{2,\lambda} > 0$ in $\overline{\Omega}$. In order to prove that $u_{2,\lambda}$ is unstable, we consider the following linearized eigenvalue problem at $u_{2,\lambda}$ with an eigenvalue γ:

$$\begin{cases} \mathcal{L}_\lambda \psi = \lambda(p-1) a u_{2,\lambda}^{p-2} \psi + \gamma \psi & \text{in } \Omega, \\ \frac{\partial \psi}{\partial \mathbf{n}} = \lambda(q-1) b u_{2,\lambda}^{q-2} \psi + \gamma \psi & \text{on } \partial\Omega, \end{cases} \tag{5.5}$$

where $\mathcal{L}_\lambda = -\Delta - \lambda m$. Let us denote by γ_1 its smallest eigenvalue and by $\psi_1 \in \mathcal{C}^{2+\alpha}(\overline{\Omega})$ an eigenfunction associated to γ_1 which is positive in $\overline{\Omega}$. We claim that $\gamma_1 < 0$. To this end, we use Picone's identity [6]. By direct computations, we have

$$\begin{aligned} \int_\Omega \left(\frac{u_{2,\lambda}}{\psi_1} \right) \sum_j \frac{\partial}{\partial x_j} \psi_1^2 \frac{\partial}{\partial x_j} \left(\frac{u_{2,\lambda}}{\psi_1} \right) &= \int_\Omega \left(\frac{u_{2,\lambda}}{\psi_1} \right) \left(-\mathcal{L}_\lambda u_{2,\lambda} \psi_1 + u_{2,\lambda} \mathcal{L}_\lambda \psi_1 \right) \\ &= \lambda(p-2) A(u_{2,\lambda}) + \gamma_1 \int_\Omega u_{2,\lambda}^2. \end{aligned}$$

On the other hand, by Green's formula we have

$$\begin{aligned} \int_\Omega \left(\frac{u_{2,\lambda}}{\psi_1} \right) \sum_j \frac{\partial}{\partial x_j} \psi_1^2 \frac{\partial}{\partial x_j} \left(\frac{u_{2,\lambda}}{\psi_1} \right) &= -\int_\Omega \psi_1^2 \left| \nabla \left(\frac{u_{2,\lambda}}{\psi_1} \right) \right|^2 + \int_{\partial\Omega} \left(\frac{u_{2,\lambda}}{\psi_1} \right) \psi_1^2 \frac{\partial}{\partial \mathbf{n}} \left(\frac{u_{2,\lambda}}{\psi_1} \right) \\ &= -\int_\Omega \psi_1^2 \left| \nabla \left(\frac{u_{2,\lambda}}{\psi_1} \right) \right|^2 + \lambda(2-q) B(u_{2,\lambda}) - \gamma_1 \int_{\partial\Omega} u_{2,\lambda}^2. \end{aligned}$$

Hence,

$$\gamma_1 = \frac{-\int_\Omega \psi_1^2 \left|\nabla\left(\frac{u_{2,\lambda}}{\psi_1}\right)\right|^2 - \lambda(p-2)A(u_{2,\lambda}) + \lambda(2-q)B(u_{2,\lambda})}{\int_\Omega u_{2,\lambda}^2 + \int_{\partial\Omega} u_{2,\lambda}^2}.$$

Since $u_{2,\lambda} \in \mathcal{N}_\lambda$, we have $\lambda B(u_{2,\lambda}) = E_\lambda(u_{2,\lambda}) - \lambda A(u_{2,\lambda})$. So, it follows that

$$\gamma_1 = \frac{-\int_\Omega \psi_1^2 \left|\nabla\left(\frac{u_{2,\lambda}}{\psi_1}\right)\right|^2 + (2-q)E_\lambda(u_{2,\lambda}) - \lambda(p-q)A(u_{2,\lambda})}{\int_\Omega u_{2,\lambda}^2 + \int_{\partial\Omega} u_{2,\lambda}^2}.$$

Since $u_{2,\lambda} \in \mathcal{N}_\lambda^-$, we have $E_\lambda(u_{2,\lambda}) < \lambda\left(\frac{p-q}{2-q}\right)A(u_{2,\lambda})$, and hence, $\gamma_1 < 0$, as desired. The proof of Theorem 1.6 is complete. □

Sketch of the proof of Remark 1.7:

Let us assume (1.14) and that w_0 is a classical positive solution of (1.12). In the same way as in the proofs of Theorems 1.3 and 1.6, we infer that, for $\lambda > 0$ sufficiently small, $u_{0,\lambda}$ is a classical positive solution of (P_λ). In order to discuss the stability of $u_{0,\lambda}$, we replace $u_{2,\lambda}$ by $u_{0,\lambda}$ in (5.5) and analyze the sign of $\gamma_1 = \gamma_1(\lambda)$. Let $\psi_1 = \psi_1(\lambda)$ be the unique positive eigenfunction associated to γ_1 and satisfying $\int_{\partial\Omega}\psi_1^2 + \int_\Omega \psi_1^2 = 1$. Setting $\xi_\lambda(x) = \lambda m(x) + \lambda(p-1)a(x)u_{0,\lambda}(x)^{p-2}$ and $\eta_\lambda(x) = \lambda(q-1)b(x)u_{0,\lambda}(x)^{q-2}$, we observe from Theorem 1.3 that $\xi_\lambda \to 0$ in $L^2(\Omega)$ and $\eta_\lambda \to (q-1)bw_0^{q-2}$ in $H^1(\Omega)$ as $\lambda \to 0^+$, where b is understood as an extension to $\mathcal{C}^{1+\alpha}(\overline{\Omega})$. By the continuity of γ_1, ψ_1 with respect to λ, we get to a limiting eigenvalue problem as $\lambda \to 0^+$, namely:

$$\begin{cases} -\Delta\psi_1(0) = \gamma_1(0)\psi_1(0) & \text{in } \Omega, \\ \frac{\partial\psi_1(0)}{\partial\mathbf{n}} = (q-1)b(x)w_0^{q-2}\psi_1(0) + \gamma_1(0)\psi_1(0) & \text{on } \partial\Omega. \end{cases}$$

By Green's formula, we have

$$\int_\Omega |\nabla\psi_1(0)|^2 - \int_{\partial\Omega}(q-1)bw_0^{q-2}\psi_1(0)^2 = \gamma_1(0). \tag{5.6}$$

Now, we claim that $\gamma_1(0) > 0$. Once this is verified, by the continuity of γ_1 we conclude that $\gamma_1(\lambda) > 0$ for $\lambda > 0$ sufficiently small, and the proof is complete. Using Green's formula again, we see that

$$0 = \int_\Omega(-\Delta w_0)\frac{\psi_1(0)^2}{w_0} = -\int_\Omega\left|\frac{\psi_1(0)}{w_0}\nabla w_0 - \nabla\psi_1(0)\right|^2 + \int_\Omega|\nabla\psi_1(0)|^2 - \int_{\partial\Omega} bw_0^{q-2}\psi_1(0)^2,$$

which combined with (5.6) yields

$$\begin{aligned}\gamma_1(0) &= (q-1)\int_\Omega\left|\frac{\psi_1(0)}{w_0}\nabla w_0 - \nabla\psi_1(0)\right|^2 + (2-q)\int_\Omega|\nabla\psi_1(0)|^2 \\ &\geq (2-q)\int_\Omega|\nabla\psi_1(0)|^2 > 0,\end{aligned}$$

since $\psi_1(0)$ is not a constant. □

Proof of Theorem 1.13:

Let

$$\mu_{1,\pm} = \inf\left\{\int_{D_\pm}\left(|\nabla u|^2 - \lambda m u^2\right);\, u \in H_0^1(D_\pm),\ \int_{D_\pm} u^2 = 1\right\}$$

be the unique positive principal eigenvalues of the Dirichlet eigenvalue problems

$$\begin{cases} -\Delta u = \lambda m u + \mu_\pm u & \text{in } D_\pm, \\ u = 0 & \text{on } \partial D_\pm, \end{cases}$$

and let $u_{1,\pm}$ denote the corresponding positive eigenfunctions in $H_0^1(D_\pm)$, respectively. By a standard regularity argument and the strong maximum principle, it follows that $u_{1,\pm} \in W^{2,r}(D_\pm)$ for any $r > N$, and $u_{1,\pm} > 0$ in $D_\pm$. Then, by Green's formula, we deduce

$$\int_{D_\pm} \nabla u_{1,\pm}\nabla v - \int_{\partial D_\pm} \frac{\partial u_{1,\pm}}{\partial \mathbf{n}} v = \lambda \int_{D_\pm} m u_{1,\pm} v + \mu_{1,\pm}\int_{D_\pm} u_{1,\pm} v \quad \text{for all } v \in \mathcal{C}^1(\overline{D_\pm}).$$

On the other hand, for any nontrivial non-negative solution $u \in H^1(\Omega)$ of (P_λ), we have

$$\int_\Omega \nabla u \nabla w - \lambda \int_\Omega m u w - \lambda \int_\Omega a u^{p-1} w - \lambda \int_{\partial\Omega} b u^{q-1} w = 0 \quad \forall\, w \in H^1(\Omega),$$

and recall that $u \in \mathcal{C}^\alpha(\overline{\Omega}) \cap W^{2,r}_{\mathrm{loc}}(\Omega)$, $r > N$, and $u > 0$ in Ω.

Now, we consider $v = u$ and $w = \widetilde{u_{1,\pm}}$, where

$$\widetilde{u_{1,\pm}} = \begin{cases} u_{1,\pm} & \text{in } D_\pm, \\ 0, & \text{otherwise,} \end{cases}$$

and then observe that

$$0 < -\int_{\partial D_\pm} \frac{\partial u_{1,\pm}}{\partial \mathbf{n}} u = \mu_{1,\pm}\int_{D_\pm} u u_{1,\pm} - \lambda \int_{D_\pm} a u^{p-1} u_{1,\pm},$$

since $u_{1,\pm} \in \mathcal{C}^1(\overline{D_\pm})$, $u_{1,\pm} > 0$ in $D_\pm$, and $\frac{\partial u_{\pm,1}}{\partial \mathbf{n}} < 0$ from the boundary point lemma (cf. [30]). Hence, it follows that $\mu_{1,+} > 0$ if $\lambda > 0$, and also that $\mu_{1,-} > 0$ if $\lambda < 0$. Since m changes sign in $D_\pm$, we have $\mu_{1,\pm} < 0$ for $|\lambda| > \Lambda$ if Λ is sufficiently large. Thus, we obtain $|\lambda| \le \Lambda$. ☐

5.1. **Bifurcating solutions.**

In this final subsection we prove Theorems 1.8 and 1.15 by a bifurcation technique. Since this technique does not require a variational structure for (P_λ), the next results hold under the condition

$$1 < q < 2 < p. \tag{5.7}$$

We use the usual orthogonal decomposition $L^2(\Omega) = \mathbb{R} \oplus V$, where

$$V = \left\{v \in L^2(\Omega) : \int_\Omega v = 0\right\},$$

and the projection $Q : L^2(\Omega) \to V$ given by

$$v = Qu = u - \frac{1}{|\Omega|}\int_\Omega u.$$

In this way we reduce the problem of finding a classical positive solution to (P_λ) under (1.14) to the following two problems

$$\begin{cases} -\Delta v + \frac{\lambda}{|\Omega|}\int_{\partial\Omega} b(x)g(t+v) = \lambda Q f(x,t+v) & \text{in } \Omega, \\ \dfrac{\partial v}{\partial \mathbf{n}} = \lambda b(x) g(t+v) & \text{on } \partial\Omega, \end{cases} \tag{5.8}$$

$$\lambda\left(\int_\Omega f(x,t+v) + \int_{\partial\Omega} bg(t+v)\right) = 0, \tag{5.9}$$

where

$$t = \frac{1}{|\Omega|}\int_\Omega u, \quad v = Qu = u - t, \quad f(x,u) = m(x)u + a(x)u^{p-1}, \quad \text{and} \quad g(u) = u^{q-1}.$$

First, to solve (5.8) in the Hölder space $\mathcal{C}^{2+\alpha}(\overline{\Omega})$, we set

$$X = \left\{ v \in \mathcal{C}^{2+\alpha}(\overline{\Omega}) : \int_\Omega v = 0 \right\}$$

and introduce the nonlinear mapping $F : \mathbb{R}\times\mathbb{R}\times X \to Z$ given by

$$F(\lambda,t,v) = \left(-\Delta v - \lambda Q f(x,t+v) + \frac{\lambda}{|\Omega|}\int_{\partial\Omega} bg(t+v),\ \frac{\partial v}{\partial \mathbf{n}} - \lambda bg(t+v)\right),$$

where

$$Z = \left\{ (\phi,\psi) \in \mathcal{C}^{\alpha}(\overline{\Omega}) \times \mathcal{C}^{1+\alpha}(\partial\Omega) : \int_\Omega \phi + \int_{\partial\Omega}\psi = 0 \right\}.$$

The Fréchet derivative of F with respect to v at $(0,c,0)$ is given by $F_v(0,c,0)v = (-\Delta v, \frac{\partial v}{\partial \mathbf{n}})$, where $c > 0$ is a constant. From Banach's closed graph theorem it follows that $F_v(0,c,0)$ is a homeomorphism. By the implicit function theorem, the set $F(\lambda,t,v) = 0$ consists exactly of an unique $\mathcal{C}^\infty$ function $v = v(\lambda,t)$ in a neighbourhood of $(\lambda,t) = (0,c)$, satisfying $v(0,c) = 0$.

Now, plugging $v(\lambda,t)$ in (5.9), we obtain the bifurcation equation

$$\lambda\left(\int_\Omega f(x,t+v(\lambda,t)) + \int_{\partial\Omega} bg(t+v(\lambda,t))\right) = 0.$$

From this equation we deduce that $\lambda = 0$ corresponds to the trivial solution $(\lambda,u) = (0,d)$ with $c-\varepsilon < d < c+\varepsilon$ for some $\varepsilon > 0$.

Hence, the study of the set of non-trivial solutions for (λ,u) close to $(0,c)$ is reduced to the consideration of the equation

$$\Phi(\lambda,t) := \int_\Omega f(x,t+v(\lambda,t)) + \int_{\partial\Omega} bg(t+v(\lambda,t)) = 0 \tag{5.10}$$

for (λ,t) close to $(0,c)$.

Recall that under the condition that $\int_\Omega m > 0 > \int_\Omega a$ and $\int_{\partial\Omega} b < 0$, the assumption $\int_{\partial\Omega} b > -\tilde{K}_1(m,a)$ made in (1.15) is equivalent to the existence of two positive zeros $c_1 < c_2$ of φ in (1.5).

Theorems 1.8 and 1.15 (1) are direct consequences of the following result.

Proposition 5.3. *Assume* (1.13), (1.14), (1.15), *and* (5.7). *Then the following two assertions hold.*

(1) (P_λ) *has two classical positive solutions* $U_{j,\lambda}$, $j = 1,2$, *for* λ *close to* 0, *given by*

$$U_{j,\lambda} = t_j(\lambda) + v(\lambda, t_j(\lambda)).$$

Here t_j is a $\mathcal{C}^1$ function of λ such that $\lambda \mapsto U_{j,\lambda} \in \mathcal{C}^{2+\alpha}(\overline{\Omega})$ is a $\mathcal{C}^1$ map, $t_j(0) = c_j$, and $v(0, c_j) = 0$, $j = 1, 2$. Moreover, $U_{1,\lambda}$ is unstable (respect. asymptotically stable), whereas $U_{2,\lambda}$ is asymptotically stable (respect. unstable) if $\lambda > 0$ (respect. $\lambda < 0$).

(2) If (P_λ) has a classical positive solution u_λ with $\lambda \neq 0$ such that $u_\lambda \to c$ in $\mathcal{C}(\overline{\Omega})$ as $\lambda \to 0$, where $c > 0$ is a constant, then $\varphi(c) = 0$.

Proof.

(1) Differentiating Φ in (5.10) with respect to t we find

$$\Phi_t(\lambda, t) = \int_\Omega f_u(x, t+v)(1+v_t) + \int_{\partial\Omega} bg'(t+v)(1+v_t).$$

From (5.8) we have

$$\begin{cases} -\Delta v_t + \frac{\lambda}{|\Omega|} \int_{\partial\Omega} bg'(t+v)(1+v_t) = \lambda Q[f_u(x, t+v)(1+v_t)] & \text{in } \Omega, \\ \dfrac{\partial v_t}{\partial \mathbf{n}} = \lambda bg'(t+v)(1+v_t) & \text{on } \partial\Omega. \end{cases}$$

Putting $\lambda = 0$ and $t = c_j$ we get

$$\begin{cases} -\Delta v_t(0, c_j) = 0 & \text{in } \Omega \\ \dfrac{\partial v_t(0, c_j)}{\partial \mathbf{n}} = 0 & \text{on } \partial\Omega. \end{cases}$$

Since $v_t(0, c_j) \in V$, we have $v_t(0, c_j) = 0$. It follows that

$$\begin{aligned} \Phi_t(0, c_j) &= \int_\Omega f_u(x, c_j) + \int_{\partial\Omega} bg'(c_j) \\ &= \int_\Omega m + (p-1)c_j^{p-2} \int_\Omega a + (q-1)c_j^{q-2} \int_{\partial\Omega} b. \end{aligned}$$

Therefore,

$$\frac{c_j^{2-q}}{q-1} \Phi_t(0, c_j) = \frac{c_j^{2-q}}{q-1} \int_\Omega m + \left(\frac{p-1}{q-1}\right) c_j^{p-q} \int_\Omega a + \int_{\partial\Omega} b. \tag{5.11}$$

Now, we claim that $\Phi_t(0, c_j) \neq 0$. Once this is verified, we end the proof of Proposition 5.3 (1) by the use of the implicit function theorem. Let $c_0 \in (c_1, c_2)$ be the global maximum point of φ. This one is given explicitly by

$$c_0 = \left(\frac{(2-q) \int_\Omega m}{(p-q)(-\int_\Omega a)} \right)^{\frac{1}{p-2}}. \tag{5.12}$$

From (5.11) and the fact that $\varphi(c_j) = 0$, we deduce that

$$\frac{c^{2-q}}{q-1} \Phi_t(0, c_j) = \left(\frac{2-q}{q-1}\right) c_j^{2-q} \int_\Omega m + \left(\frac{p-q}{q-1}\right) c_j^{p-q} \int_\Omega a.$$

It follows from (5.12) that

$$\begin{aligned} \frac{1}{q-1} \Phi_t(0, c_j) &= \left(\frac{2-q}{q-1}\right) \int_\Omega m + \left(\frac{p-q}{q-1}\right) c_j^{p-2} \int_\Omega a \\ &\begin{cases} > \left(\frac{2-q}{q-1}\right) \int_\Omega m + \left(\frac{p-q}{q-1}\right) c_0^{p-2} \int_\Omega a = 0, & \text{for } j = 1, \\ < \left(\frac{2-q}{q-1}\right) \int_\Omega m + \left(\frac{p-q}{q-1}\right) c_0^{p-2} \int_\Omega a = 0, & \text{for } j = 2. \end{cases} \end{aligned}$$

The conclusion follows.

We prove now the stability results of $U_{j,\lambda}$. We recall from (5.5) the linearized eigenvalue problem at $U_{j,\lambda}$:

$$\begin{cases} -\Delta\psi = \lambda m\psi + \lambda(p-1)U_{j,\lambda}^{p-2}\psi + \gamma\psi & \text{in } \Omega, \\ \frac{\partial\psi}{\partial\mathbf{n}} = \lambda(q-1)U_{j,\lambda}^{q-2}\psi + \gamma\psi & \text{on } \partial\Omega. \end{cases} \tag{5.13}$$

Let $\gamma_1 = \gamma_1(\lambda)$ be the smallest positive eigenvalue of this problem and $\psi_1 = \psi_1(\lambda)$ be the unique positive eigenfunction associated to γ_1, satisfying $\int_\Omega \psi_1^2 + \int_{\partial\Omega} \psi_1^2 = 1$. It is easy to see that $\gamma_1(0) = 0$ and $\psi_1(0) = \left(\frac{1}{|\Omega|+|\partial\Omega|}\right)^{1/2}$. We differentiate (5.13) with respect to λ and let $\lambda = 0$ to obtain

$$\begin{cases} -\Delta\psi_1'(0) = m\psi_1(0) + (p-1)aU_{j,0}^{p-2}\psi_1(0) + \gamma_1'(0)\psi_1(0) & \text{in } \Omega, \\ \frac{\partial\psi_1'(0)}{\partial\mathbf{n}} = (q-1)bU_{j,0}^{q-2}\psi_1(0) + \gamma_1'(0)\psi_1(0) & \text{on } \partial\Omega. \end{cases}$$

By Green's formula, we have

$$\int_\Omega \Big(m\psi_1(0) + (p-1)aU_{j,0}^{p-2}\psi_1(0) + \gamma_1'(0)\psi_1(0) \Big) \\ + \int_{\partial\Omega} \Big((q-1)bU_{j,0}^{q-2}\psi_1(0) + \gamma_1'(0)\psi_1(0) \Big) = 0.$$

Since $U_{j,0} = c_j$ and $\varphi(c_j) = 0$ from (1.5), we deduce

$$\gamma_1'(0) = -\frac{c_j^{q-2}\left\{(2-q)c_j^{2-q}\int_\Omega m + (p-q)c_j^{p-q}\int_\Omega a\right\}}{|\Omega| + |\partial\Omega|}. \tag{5.14}$$

By a direct computation, we see that

$$c_j\varphi'(c_j) = (2-q)c_j^{2-q}\int_\Omega m + (p-q)c_j^{p-q}\int_\Omega a.$$

Since $\varphi(c_1) > 0 > \varphi_1(c_2)$, we deduce from (5.14) that $\gamma_1'(0) < 0$ for $j = 1$ and $\gamma_1'(0) > 0$ for $j = 2$, which combined with $\gamma_1(0) = 0$ provides the desired conclusion.

(2) Let u be a classical positive solution of (P_λ) with $\lambda \neq 0$. By Green's formula it follows that

$$\int_\Omega -\Delta u = -\lambda\int_{\partial\Omega} bu^{q-1}.$$

Hence we have

$$\lambda\left(\int_\Omega (mu + au^{p-1}) + \int_{\partial\Omega} bu^{q-1}\right) = 0,$$

i.e.

$$\int_\Omega (mu + au^{p-1}) + \int_{\partial\Omega} bu^{q-1} = 0.$$

Since $u \to c$ in $\mathcal{C}(\overline{\Omega})$, where c is a positive constant, we obtain the desired conclusion.

□

Next, we recall that under the conditions $\int_\Omega m > 0 > \int_\Omega a$ and $\int_{\partial\Omega} b < 0$, the assumption $\int_{\partial\Omega} b = -\tilde{K}_1(m,a)$ is equivalent to the existence of a unique positive zero c_0 of φ, given by (5.12). In this case, $\varphi(c_0) = \varphi'(c_0) = 0$, and c_0 is the global maximum point of φ.

Lemma 5.4. *Assume* (1.14), (5.7), $\int_\Omega m > 0 > \int_\Omega a$, *and* $\int_{\partial\Omega} b = -\tilde{K}_1(m,a)$. *Then, for* Φ *defined in* (5.10), *we have the following:*

(1) $\Phi(0, c_0) = 0$.
(2) $\Phi_t(0, c_0) = 0$.
(3) $\Phi_{tt}(0, c_0) = -(2-q)(p-2)c_0^{-1} \int_\Omega m < 0$.
(4) $\Phi_\lambda(0, c_0) = \int_\Omega m v_\lambda(0, c_0) + (p-1)c_0^{p-2} \int_\Omega a v_\lambda(0, c_0) + (q-1)c_0^{q-2} \int_{\partial\Omega} b v_\lambda(0, c_0)$.

In particular, if $a = -km$ for some positive constant k, then

$$\Phi_\lambda(0, c_0) = (q-1)c_0^{-1} \int_\Omega |\nabla v_\lambda(0, c_0)|^2 > 0. \tag{5.15}$$

Proof.

(1) It is straightforward from

$$\Phi(0, c_0) = \int_\Omega (mc_0 + ac_0^{p-1}) + \int_{\partial\Omega} bc_0^{q-1} = c_0^{q-1}\varphi(c_0) = 0.$$

(2) We differentiate Φ with respect to t to get

$$\Phi_t = \int_\Omega f_u(x, t+v)(1+v_t) + \int_{\partial\Omega} bg'(t+v)(1+v_t).$$

Thus

$$\Phi_t(0, c_0) = \int_\Omega f_u(x, c_0)(1 + v_t(0, c_0)) + \int_{\partial\Omega} bg'(c_0)(1 + v_t(0, c_0)). \tag{5.16}$$

Let us show how we derive $v_t(0, c_0)$ from (5.8). Differentiating (5.8) (with $v = v(\lambda, t)$) with respect to t, we obtain

$$\begin{cases} -\Delta v_t + \frac{\lambda}{|\Omega|} \int_{\partial\Omega} bg'(t+v)(1+v_t) = \lambda Q[f_u(x, t+v)(1+v_t)] & \text{in } \Omega, \\ \dfrac{\partial v_t}{\partial \mathbf{n}} = \lambda bg'(t+v)(1+v_t) & \text{on } \partial\Omega. \end{cases}$$

Taking $(\lambda, t) = (0, c_0)$ we get

$$\begin{cases} -\Delta v_t(0, c_0) = 0 & \text{in } \Omega, \\ \dfrac{\partial v_t(0, c_0)}{\partial \mathbf{n}} = 0 & \text{on } \partial\Omega. \end{cases}$$

Since $v_t(0, c_0) \in V$, i.e. $\int_\Omega v_t(0, c_0) = 0$, we have

$$v_t(0, c_0) = 0. \tag{5.17}$$

Hence, from (5.16) and $\varphi(c_0) = \varphi'(c_0) = 0$, it follows that

$$\begin{aligned} \Phi_t(0, c_0) &= \int_\Omega (m + (p-1)ac_0^{p-2}) + \int_{\partial\Omega} b(q-1)c_0^{q-2} \\ &= \int_\Omega (m + (p-1)ac_0^{p-2}) + (q-1)c_0^{q-2} \int_{\partial\Omega} b \\ &= \int_\Omega (m + (p-1)ac_0^{p-2}) + (q-1)c_0^{q-2} \left(-c_0^{p-q} \int_\Omega a - c_0^{2-q} \int_\Omega m \right) \\ &= (2-q) \int_\Omega m + (p-q)c_0^{p-2} \int_\Omega a = 0. \end{aligned}$$

(3) Differentiating Φ once more with respect to t, we have

$$\Phi_{tt}(0,c_0) = \int_\Omega \{f_{uu}(x,c_0)(1+v_t(0,c_0))^2 + f_u(x,c_0)v_{tt}(0,c_0)\} \\ + \int_{\partial\Omega} b\{g''(c_0)(1+v_t(0,c_0))^2 + g'(c_0)v_{tt}(0,c_0)\}.$$

In the same way as $v_t(0,c_0)=0$, we get $v_{tt}(0,c_0)=0$ from (5.8). It follows that

$$\begin{aligned}
\Phi_{tt}(0,c_0) &= (p-1)(p-2)c_0^{p-3}\int_\Omega a + (q-1)(q-2)c_0^{q-3}\int_{\partial\Omega} b \\
&= (p-1)(p-2)c_0^{p-3}\int_\Omega a + (q-1)(q-2)c_0^{q-3}\left(-c_0^{p-q}\int_\Omega a - c_0^{2-q}\int_\Omega m\right) \\
&= \{(p-1)(p-2)-(q-1)(q-2)\}c_0^{p-3}\int_\Omega a - (q-1)(q-2)c_0^{-1}\int_\Omega m \\
&= c_0^{-1}\left(\{(p-1)(p-2)-(q-1)(q-2)\}c_0^{p-2}\int_\Omega a - (q-1)(q-2)\int_\Omega m\right) \\
&= c_0^{-1}\left(\{(p-1)(p-2)-(q-1)(q-2)\}\frac{(2-q)\int_\Omega m}{(p-q)(-\int_\Omega a)}\int_\Omega a - (q-1)(q-2)\int_\Omega m\right) \\
&= c_0^{-1}\int_\Omega m\left(\frac{2-q}{p-q}\right)\{(q-1)(q-2)+(p-1)(p-2)-(q-1)(p-q)\} \\
&= c_0^{-1}\int_\Omega m\left(\frac{2-q}{p-q}\right)(p-2)(q-p) = -(2-q)(p-2)c_0^{-1}\int_\Omega m < 0,
\end{aligned}$$

where we have used again that $\varphi(c_0)=\varphi'(c_0)=0$.

(4) From the formula

$$\Phi_\lambda = \int_\Omega f_u(x,t+v)v_\lambda + \int_{\partial\Omega} bg'(t+v)v_\lambda,$$

it follows that

$$\begin{aligned}
\Phi_\lambda(0,c_0) &= \int_\Omega f_u(x,c_0)v_\lambda(0,c_0) + \int_{\partial\Omega} bg'(c_0)v_\lambda(0,c_0) \\
&= \int_\Omega (m+(p-1)ac_0^{p-2})v_\lambda(0,c_0) + \int_{\partial\Omega} b(q-1)c_0^{q-2}v_\lambda(0,c_0) \\
&= \int_\Omega mv_\lambda(0,c_0) + (p-1)c_0^{p-2}\int_\Omega av_\lambda(0,c_0) + (q-1)c_0^{q-2}\int_{\partial\Omega} bv_\lambda(0,c_0).
\end{aligned}$$

(5) From (5.8) we get

$$\begin{cases} -\Delta v_\lambda + \frac{1}{|\Omega|}\int_{\partial\Omega} b[g(t+v)+\lambda g'(t+v)v_\lambda] = Q[f(x,t+v)] + \lambda Q[f_u(x,t+v)v_\lambda] & \text{in } \Omega, \\ \dfrac{\partial v_\lambda}{\partial \mathbf{n}} = b[g(t+v)+\lambda g'(t+v)v_\lambda] & \text{on } \partial\Omega. \end{cases}$$

Put $\lambda=0$, $t=c_0$, and $v(0,c_0)=0$, to obtain

$$\begin{cases} -\Delta v_\lambda(0,c_0) + \frac{1}{|\Omega|}\int_{\partial\Omega} bg(c_0) = Q[f(x,c_0)] & \text{in } \Omega, \\ \dfrac{\partial v_\lambda(0,c_0)}{\partial \mathbf{n}} = bg(c_0) & \text{on } \partial\Omega, \end{cases}$$

where

$$Q[f(x,c_0)] = mc_0 + ac_0^{p-1} - \frac{1}{|\Omega|}\int_\Omega (mc_0 + ac_0^{p-1}).$$

It follows that

$$\begin{cases} -\Delta v_\lambda(0,c_0) + \frac{1}{|\Omega|}\{\int_\Omega (mc_0 + ac_0^{p-1}) + \int_{\partial\Omega} bc_0^{q-1}\} = mc_0 + ac_0^{p-1} & \text{in } \Omega, \\ \dfrac{\partial v_\lambda(0,c_0)}{\partial \mathbf{n}} = bc_0^{q-1} & \text{on } \partial\Omega, \end{cases}$$

and consequently

$$\begin{cases} -\Delta v_\lambda(0,c_0) = mc_0 + ac_0^{p-1} & \text{in } \Omega, \\ \dfrac{\partial v_\lambda(0,c_0)}{\partial \mathbf{n}} = bc_0^{q-1} & \text{on } \partial\Omega, \end{cases}$$

since $\varphi(c_0) = 0$. Hence

$$\int_\Omega |\nabla v_\lambda(0,c_0)|^2 - c_0^{q-1}\int_{\partial\Omega} bv_\lambda(0,c_0) = \int_\Omega (mc_0 + ac_0^{p-1})v_\lambda(0,c_0).$$

From (4) we get

$$\begin{aligned} c_0\Phi_\lambda(0,c_0) &= c_0\int_\Omega mv_\lambda(0,c_0) + (p-1)c_0^{p-1}\int_\Omega av_\lambda(0,c_0) + (q-1)c_0^{q-1}\int_{\partial\Omega} bv_\lambda(0,c_0) \\ &= (q-1)\int_\Omega |\nabla v_\lambda(0,c_0)|^2 + \int_\Omega \Big\{(2-q)c_0m + (p-q)c_0^{p-1}a\Big\} v_\lambda(0,c_0). \end{aligned}$$

Since k is a positive constant and $a = -km$, we have

$$\begin{aligned} (2-q)c_0m + (p-q)c_0^{p-1}a &= mc_0\Big\{(2-q) - (p-q)c_0^{p-2}k\Big\} \\ &= mc_0\left\{(2-q) - (p-q)\frac{(2-q)\int_\Omega m}{(p-q)(-\int_\Omega a)}k\right\} \\ &= 0. \end{aligned}$$

Therefore

$$c_0\Phi_\lambda(0,c_0) = (q-1)\int_\Omega |\nabla v_\lambda(0,c_0)|^2.$$

Moreover, since $b \not\equiv 0$, $v_\lambda(0,c_0)$ is not a constant, so that $\int_\Omega |\nabla v_\lambda(0,c_0)|^2 > 0$.

The proof of Lemma 5.4 is now complete. □

Theorem 1.15 (2) is then a direct consequence of the following result:

Proposition 5.5. *Assume* (1.14), (5.7), $a = -km$ *for some positive constant* k *and*

$$\int_\Omega m > 0 > \int_{\partial\Omega} b = -\tilde{K}_1(m,a).$$

Then there exists a constant $\varepsilon > 0$ *and a* $\mathcal{C}^3$ *function* $\lambda : (c_0 - \varepsilon, c_0 + \varepsilon) \to \mathbb{R}$ *satisfying* $\lambda(c_0) = \lambda'(c_0) = 0$ *and* $\lambda''(c_0) > 0$, *such that the set*

$$\{(\lambda(t),\ t + v(\lambda(t),t)) : t \in (c_0 - \varepsilon, c_0 + \varepsilon)\}$$

is contained in the positive solutions set of (P_λ). *Moreover, the positive solution* $t + v(\lambda(t),t)$ *of* $(P_{\lambda(t)})$ *is asymptotically stable for* $c_0 < t < c_0 + \varepsilon$ *and unstable for* $c_0 - \varepsilon < t < c_0$.

Remark 5.6. From (5.12) note that if $a = -km$ then $c_0 = c_0(k) = \left(\frac{2-q}{(p-q)k}\right)^{\frac{1}{p-2}}$. It follows that $k \mapsto c_0(k)$ is decreasing, $\lim_{k\to 0^+} c_0(k) = \infty$ and $\lim_{k\to\infty} c_0(k) = 0$.

Proof. By (5.15) and the implicit function theorem, we deduce that there exists a $\mathcal{C}^3$ function $t \mapsto \lambda(t)$ such that

$$\Phi(\lambda, t) = 0 \ \text{ for } (\lambda, t) \text{ close to } (0, c_0) \Longleftrightarrow (\lambda, t) = (\lambda(t), t) \ \text{ for } t \text{ close to } c_0.$$

From Lemma 5.4 (1) we have $\lambda(c_0) = 0$, whereas from Lemma 5.4 (2), (3), and (5.15), we have

$$\lambda'(c_0) = -\frac{\Phi_t(0, c_0)}{\Phi_\lambda(0, c_0)} = 0, \quad \lambda''(c_0) = -\frac{\Phi_{tt}(0, c_0)}{\Phi_\lambda(0, c_0)} > 0,$$

as desired.

We prove now the stability result. Recall c_0 is the unique zero of φ, given by (1.5), as well as its global maximum point. So we have

$$c_0^{q-2}\varphi(c_0) = \int_\Omega m + c_0^{p-2}\int_\Omega a + c_0^{q-2}\int_{\partial\Omega} b = 0, \tag{5.18}$$

$$c_0^{q-1}\varphi'(c_0) = (2-q)\int_\Omega m + (p-q)c_0^{p-2}\int_\Omega a = 0. \tag{5.19}$$

We let $w(t) := t + v(\lambda(t), t)$, and consider the stability of $(\lambda(t), w(t))$, $|t - c_0| < \varepsilon$. To this end, we study the linearized eigenvalue problem at $(\lambda(t), w(t))$, which is given by

$$\begin{cases} -\Delta\psi = \lambda m\psi + \lambda a(p-1)w^{p-2}\psi + \gamma\psi & \text{in } \Omega, \\ \partial_{\mathbf{n}}\psi = \lambda b(q-1)w^{q-2}\psi + \gamma\psi & \text{on } \partial\Omega. \end{cases} \tag{5.20}$$

Let $\gamma_1 = \gamma_1(t)$ be its smallest eigenvalue, which is simple, and $\psi_1 = \psi_1(t)$ be the positive eigenfunction associated to γ_1 satisfying $\int_\Omega \psi_1^2 + \int_{\partial\Omega} \psi_1^2 = 1$.

First, we claim that

$$\gamma_1(c_0) = 0, \quad \text{and} \ \ \psi_1(c_0) \equiv \left(\frac{1}{|\Omega| + |\partial\Omega|}\right)^{1/2}. \tag{5.21}$$

Indeed, putting $t = c_0$ in (5.20), we have

$$\begin{cases} -\Delta\psi_1(c_0) = \gamma_1(c_0)\psi_1(c_0) & \text{in } \Omega, \\ \partial_{\mathbf{n}}\psi_1 = \gamma_1(c_0)\psi_1(c_0) & \text{on } \partial\Omega. \end{cases}$$

By uniqueness, it follows that $\gamma_1(c_0) = 0$ and

$$\psi_1(c_0) \equiv \left(\frac{1}{|\Omega| + |\partial\Omega|}\right)^{1/2},$$

as claimed.

Second, we show that

$$\gamma_1'(c_0) = 0, \quad \text{and} \ \ \psi_1'(c_0) = 0. \tag{5.22}$$

To this end, we differentiate (5.20) with respect to t (with $\gamma = \gamma_1$ and $\psi = \psi_1$) to obtain

$$\begin{cases} -\Delta\psi_1' = m\left(\lambda'\psi_1 + \lambda\psi_1'\right) + (p-1)a\left(\lambda' w^{p-2}\psi_1 + \lambda(p-2)w^{p-3}w'\psi_1 + \lambda w^{p-2}\psi_1'\right) & \\ \qquad\qquad + \gamma_1'\psi_1 + \gamma_1\psi_1' & \text{in } \Omega, \\ \partial_{\mathbf{n}}\psi_1' = b(q-1)\left(\lambda' w^{q-2}\psi_1 + \lambda(q-2)w^{q-3}w'\psi_1 + \lambda w^{q-2}\psi_1'\right) + \gamma_1'\psi_1 + \gamma_1\psi_1' & \text{on } \partial\Omega. \end{cases} \tag{5.23}$$

Taking $t = c_0$ in (5.23) we get

$$\begin{cases} -\Delta\psi_1'(c_0) = \gamma_1'(c_0)\psi_1(c_0) & \text{in } \Omega, \\ \partial_{\mathbf{n}}\psi_1'(c_0) = \gamma_1'(c_0)\psi_1(c_0) & \text{on } \partial\Omega. \end{cases}$$

By Green's formula, we deduce that

$$\int_\Omega (\gamma_1'(c_0)\psi_1(c_0)) + \int_{\partial\Omega} (\gamma_1'(c_0)\psi_1(c_0)) = 0,$$

so that $\gamma_1'(c_0)\psi_1(c_0)(|\Omega| + |\partial\Omega|) = 0$. Hence, we have $\gamma_1'(c_0) = 0$, and consequently $\psi_1'(c_0)$ is a constant. On the other hand, differentiating $\int_\Omega \psi_1^2 + \int_{\partial\Omega} \psi_1^2 = 1$ with respect to t we obtain

$$\int_\Omega \psi_1\psi_1' + \int_{\partial\Omega} \psi_1\psi_1' = 0. \tag{5.24}$$

Thus we have $\psi_1(c_0)\psi_1'(c_0)(|\Omega| + |\partial\Omega|) = 0$, which implies $\psi_1'(c_0) = 0$, as desired.

Third, we verify that

$$\gamma_1''(c_0) = 0. \tag{5.25}$$

Differentiating (5.23) with respect to t, we obtain

$$\begin{cases} -\Delta\psi_1'' = m\left(\lambda''\psi_1 + 2\lambda'\psi_1' + \lambda\psi''\right) + (p-1)a\left(\lambda'' w^{p-2}\psi_1 + \lambda'(w^{p-2}\psi_1)'\right) & \\ \qquad + \left[\lambda a(p-1)\left((p-2)w^{p-3}w'\psi_1 + w^{p-2}\psi_1'\right)\right]' + \gamma_1''\psi_1 + 2\gamma_1'\psi_1' + \gamma_1\psi_1'' & \text{in } \Omega, \\ \partial_{\mathbf{n}}\psi_1'' = b(q-1)\left[\lambda'' w^{q-2}\psi_1 + \lambda'(w^{q-2}\psi_1)'\right] + \left[\lambda b(q-1)\left((q-2)w^{q-3}w'\psi_1 + w^{q-2}\psi_1'\right)\right]' & \\ \qquad + \gamma_1''\psi_1 + 2\gamma_1'\psi_1' + \gamma_1\psi_1'' & \text{on } \partial\Omega. \end{cases} \tag{5.26}$$

Taking $t = c_0$ we get

$$\begin{cases} -\Delta\psi_1''(c_0) = \lambda''(c_0)m\psi_1(c_0) + \lambda''(c_0)a(p-1)(w(c_0))^{p-2}\psi_1(c_0) + \gamma_1''(c_0)\psi_1(c_0) & \text{in } \Omega, \\ \partial_{\mathbf{n}}\psi_1''(c_0) = \lambda''(c_0)b(q-1)(w(c_0))^{q-2}\psi_1(c_0) + \gamma_1''(c_0)\psi_1(c_0) & \text{on } \partial\Omega. \end{cases}$$

Since $\psi_1(c_0)$ is a positive constant, it follows by Green's formula that

$$\lambda''(c_0)\left(\int_\Omega m + (p-1)\int_\Omega a(w(c_0))^{p-2} + (q-1)\int_{\partial\Omega} b(w(c_0))^{q-2}\right) + \gamma_1''(c_0)(|\Omega| + |\partial\Omega|) = 0.$$

Note that $w(c_0) = c_0$, which combined with (5.18) and (5.19) implies

$$\int_\Omega m + (p-1)\int_\Omega a(w(c_0))^{p-2} + (q-1)\int_{\partial\Omega} b(w(c_0))^{q-2} = 0, \tag{5.27}$$

and (5.25) follows.

Finally, we verify that

$$\gamma_1'''(c_0) > 0. \tag{5.28}$$

We differentiate once more (5.26) with respect to t and take $t = c_0$. Since $\lambda(c_0) = \lambda'(c_0) = 0$ and $\psi_1'(c_0) = 0$ we deduce that

$$\begin{cases} -\Delta\psi_1'''(c_0) = \lambda'''(c_0)m\psi_1(c_0) + \lambda'''(c_0)a(p-1)(w(c_0))^{p-2}\psi_1(c_0) & \\ \qquad + 3\lambda''(c_0)a(p-1)(p-2)(w(c_0))^{p-3}w'(c_0)\psi_1(c_0) + \gamma_1'''(c_0)\psi_1(c_0) & \text{in } \Omega, \\ \partial_{\mathbf{n}}\psi_1'''(c_0) = \lambda'''(c_0)b(q-1)(w(c_0))^{q-2}\psi_1(c_0) + 3\lambda''(c_0)b(q-1)(q-2)(w(c_0))^{q-3}w'(c_0)\psi_1(c_0) & \\ \qquad + \gamma_1'''(c_0)\psi_1(c_0) & \text{on } \partial\Omega. \end{cases}$$

Since $w(c_0) = c_0$ and $\psi_1(c_0)$ is a positive constant, by Green's formula, we deduce that

$$\begin{aligned} 0 = \gamma_1'''(c_0)(|\Omega| + |\partial\Omega|) + \lambda'''(c_0)\left\{\int_\Omega \left(m + (p-1)ac_0^{p-2}\right) + (q-1)c_0^{q-2}\int_{\partial\Omega} b\right\} \\ + 3\lambda''(c_0)\left\{(p-1)(p-2)c_0^{p-3}\int_\Omega aw'(c_0) + (q-1)(q-2)c_0^{q-3}\int_{\partial\Omega} bw'(c_0)\right\}. \end{aligned}$$

Using (5.27), we see that the second term on the right hand side vanishes. In order to deal with the third term, we consider $w'(c_0)$. Note that, from the definition of w, we have $w'(t) = 1 + v_\lambda \lambda' + v_t$. Since $\lambda'(c_0) = 0$, we have then $w'(c_0) = 1 + v_t(0, c_0) = 1$, where we have used (5.17). From $a = -km$, it follows that

$$\gamma_1'''(c_0)(|\Omega| + |\partial\Omega|) = 3\lambda''(c_0)\left\{k(p-1)(p-2)c_0^{p-3}\int_\Omega m + (q-1)(2-q)c_0^{q-3}\int_{\partial\Omega} b\right\}.$$

Now, we recall the assumption

$$\int_{\partial\Omega} b = -\tilde{K}_1(m, -km), \tag{5.29}$$

which yields $\int_{\partial\Omega} b = -\tilde{C}_{pq} k^{\frac{q-2}{p-2}} \int_\Omega m$, where

$$\tilde{C}_{pq} = \left(\frac{q}{2}\left(\frac{p}{2}\right)^{\frac{2-q}{p-2}}\right)^{-1} \frac{q(p-2)}{2(p-q)}\left(\frac{p(2-q)}{2(p-q)}\right)^{\frac{2-q}{p-2}} = \frac{p-2}{p-q}\left(\frac{2-q}{p-q}\right)^{\frac{2-q}{p-2}}.$$

Moreover, from (5.19) we also have

$$c_0^{p-q} = k^{\frac{q-p}{p-2}}\left(\frac{2-q}{p-q}\right)^{\frac{p-q}{p-2}}.$$

Thus, from (5.29) we deduce that

$$\begin{aligned}
\gamma_1'''(c_0)(|\Omega| + |\partial\Omega|) &= 3\lambda''(c_0)c_0^{q-3}\left\{(p-1)(p-2)c_0^{p-q}\int_\Omega m + (q-1)(2-q)\int_{\partial\Omega} b\right\} \\
&= 3\lambda''(c_0)c_0^{q-3}\left(\int_\Omega m\right)k^{\frac{q-2}{p-2}}\left(\frac{2-q}{p-q}\right)^{\frac{2-q}{p-2}}(p-2)(2-q).
\end{aligned}$$

From $\lambda''(c_0) > 0$ and $\int_\Omega m > 0$, we infer (5.28).

Summing up, from (5.21), (5.22), (5.25), and (5.28), the desired conclusion follows.

The proof of Proposition 5.5 is now complete. □

Appendix A. A comparison principle for mixed Dirichlet and Neumann nonlinear boundary conditions

In this Appendix, we provide a variant of the comparison principle proved by Ambrosetti, Brezis and Cerami [3, Lemma 3.3] to mixed Dirichlet and Neumann nonlinear boundary conditions. We consider the general boundary value problem with mixed nonlinear boundary conditions

$$\begin{cases} -\Delta u = f(x, u) & \text{in } D, \\ \frac{\partial u}{\partial \mathbf{n}} = g(x, u) & \text{on } \Gamma_1, \\ u = 0 & \text{on } \Gamma_0, \end{cases} \tag{A.1}$$

where:

- D is a bounded domain of $\mathbb{R}^N$ with smooth boundary ∂D.
- $\Gamma_0, \Gamma_1 \subset \partial D$ are disjoint, open, and smooth $(N-1)$ dimensional surfaces of ∂D.
- $\overline{\Gamma_0}, \overline{\Gamma_1}$ are compact manifolds with $(N-2)$ dimensional closed boundary $\gamma = \overline{\Gamma_0} \cap \overline{\Gamma_1}$ such that $\partial D = \Gamma_0 \cup \gamma \cup \Gamma_1$.
- $f : \overline{\Omega} \times [0, \infty) \to \mathbb{R}$ and $g : \overline{\Gamma_1} \times [0, \infty) \to \mathbb{R}$ are continuous.

Proposition A.1. *Under the above conditions, assume that for every* $x \in D$, $t \mapsto \frac{f(x,t)}{t}$ *is decreasing in* $(0,\infty)$, *and for every* $x \in \Gamma_1$, $t \mapsto \frac{g(x,t)}{t}$ *is non-increasing in* $(0,\infty)$. *Let* $u, v \in H^1(D) \cap \mathcal{C}(\overline{D})$ *be non-negative functions satisfying* $u = 0 \leq v$ *on* Γ_0, *and*

$$\int_D \nabla u \nabla \varphi - \int_D f(x,u)\varphi - \int_{\Gamma_1} g(x,u)\varphi \leq \ 0, \quad \forall \varphi \in H^1_{\Gamma_0}(D) \ \text{ such that } \ \varphi \geq 0,$$

$$\int_D \nabla v \nabla \varphi - \int_D f(x,v)\varphi - \int_{\Gamma_1} g(x,v)\varphi \geq \ 0, \quad \forall \varphi \in H^1_{\Gamma_0}(D) \ \text{ such that } \ \varphi \geq 0.$$

If $u, v > 0$ *in* D, *then* $u \leq v$ *in* $\overline{D}$.

Proof. Let $\theta : \mathbb{R} \to \mathbb{R}$, be a nonnegative nondecreasing smooth function such that $\theta(t) = 0$ for $t \leq 0$ and $\theta(t) = 1$ for $t \geq 1$. For $\varepsilon > 0$ we set $\theta_\varepsilon(t) = \theta(t/\varepsilon)$. Since $u - v \leq 0$ on Γ_0, we have $v\theta_\varepsilon(u-v) \in H^1_{\Gamma_0}(D)$, so that

$$\int_D \nabla u \nabla (v\theta_\varepsilon(u-v)) - \int_D f(x,u) v\theta_\varepsilon(u-v) - \int_{\Gamma_1} g(x,u) v\theta_\varepsilon(u-v) \leq 0. \tag{A.2}$$

Likewise, since $u\theta_\varepsilon(u-v) \in H^1_{\Gamma_0}(D)$, we have

$$\int_D \nabla v \nabla (u\theta_\varepsilon(u-v)) - \int_D f(x,v) u\theta_\varepsilon(u-v) - \int_{\Gamma_1} g(x,v) u\theta_\varepsilon(u-v) \geq 0. \tag{A.3}$$

Let $\Gamma_1^+ = \{x \in \Gamma_1 : u, v > 0\}$. Since $t \mapsto \frac{g(x,t)}{t}$ is non-increasing in $(0,\infty)$, we have $g(x,0) \geq 0$, which combined with (A.2) and (A.3) yields

$$\begin{aligned}
&\int_D u\theta_\varepsilon'(u-v)\nabla v(\nabla u - \nabla v) - \int_D v\theta_\varepsilon'(u-v)\nabla u(\nabla u - \nabla v) \\
&\geq \int_D uv\left(\frac{f(x,v)}{v} - \frac{f(x,u)}{u}\right)\theta_\varepsilon(u-v) + \int_{\Gamma_1^+} uv\left(\frac{g(x,v)}{v} - \frac{g(x,u)}{u}\right)\theta_\varepsilon(u-v) \\
&\geq \int_D uv\left(\frac{f(x,v)}{v} - \frac{f(x,u)}{u}\right)\theta_\varepsilon(u-v).
\end{aligned}$$

From $-\int_D u\theta_\varepsilon'(u-v)|\nabla(u-v)|^2 \leq 0$, it follows that

$$\int_D (u-v)\theta_\varepsilon'(u-v)\nabla u \nabla(u-v) \geq \int_D uv\left(\frac{f(x,v)}{v} - \frac{f(x,u)}{u}\right)\theta_\varepsilon(u-v). \tag{A.4}$$

Now, we introduce $\gamma_\varepsilon(t) = \int_0^t s\theta_\varepsilon'(s)ds$ for $t \in \mathbb{R}$. We have then $0 \leq \gamma_\varepsilon(t) \leq \varepsilon$, $t \in \mathbb{R}$. Note that $\nabla(\gamma_\varepsilon(u-v)) = (u-v)\theta_\varepsilon'(u-v)\nabla(u-v)$. Hence, from (A.4) we deduce that

$$\int_D \nabla u \nabla(\gamma_\varepsilon(u-v)) \geq \int_D uv\left(\frac{f(x,v)}{v} - \frac{f(x,u)}{u}\right)\theta_\varepsilon(u-v).$$

Now, since $\gamma_\varepsilon(u-v) \in H^1_{\Gamma_0}(D)$ and $\gamma_\varepsilon(u-v) \geq 0$, we note that

$$\int_D \nabla u \nabla(\gamma_\varepsilon(u-v)) - \int_D f(x,u)\gamma_\varepsilon(u-v) - \int_{\Gamma_1} g(x,u)\gamma_\varepsilon(u-v) \leq 0,$$

and combining the two latter assertions, we get

$$\int_D f(x,u)\gamma_\varepsilon(u-v) + \int_{\Gamma_1} g(x,u)\gamma_\varepsilon(u-v) \geq \int_D uv\left(\frac{f(x,v)}{v} - \frac{f(x,u)}{u}\right)\theta_\varepsilon(u-v).$$

Since $\gamma_\varepsilon(t) \leq \varepsilon$, there exists a constant $C > 0$ such that

$$C\varepsilon \geq \int_D uv\left(\frac{f(x,v)}{v} - \frac{f(x,u)}{u}\right)\theta_\varepsilon(u-v). \tag{A.5}$$

Since $t \mapsto \frac{f(x,t)}{t}$ is decreasing in $(0,\infty)$, we use Fatou's lemma to deduce from (A.5) that

$$\int_D \liminf_{\varepsilon \to 0^+} uv \left(\frac{f(x,v)}{v} - \frac{f(x,u)}{u} \right) \theta_\varepsilon(u-v) \leq 0.$$

Note that

$$\lim_{\varepsilon \to 0^+} \theta_\varepsilon(u-v) = \begin{cases} 1, & u > v, \\ 0, & u \leq v, \end{cases}$$

so that

$$\int_{\{u>v\}} uv \left(\frac{f(x,v)}{v} - \frac{f(x,u)}{u} \right) \leq 0.$$

Using again that $t \mapsto \frac{f(x,t)}{t}$ is decreasing in $(0,\infty)$, we conclude $|\{u > v\}| = 0$, which implies $u \leq v$ a.e. in D. By continuity, the desired conclusion follows. □

Appendix B. Positivity of nontrivial non-negative weak solutions in the one-dimensional case

In this Appendix, we show the positivity of nontrivial non-negative weak solutions for the one-dimensional case of (P_λ). We take $\Omega = I = (0,1)$ and show that under some regularity assumptions on m and a a nontrivial non-negative solution satisfies $u > 0$ on $\overline{I}$. More precisely, we consider nontrivial non-negative weak solutions of the problem

$$\begin{cases} -u'' = \lambda(m(x)u + a(x)u^{p-1}) \quad \text{in } I, \\ -u'(0) = \lambda b_0 u(0)^{q-1}, \\ u'(1) = \lambda b_1 u(1)^{q-1}, \end{cases} \tag{B.1}$$

where $1 < q < 2 < p$, $m, a \in \mathcal{C}^1(\overline{I})$, and $b_0, b_1 \in \mathbb{R}$. A non-negative function $u \in H^1(I)$ is a non-negative weak solution of (B.1) if it satisfies

$$\int_I u'\phi' = \lambda \left(b_0 u(0)^{q-1}\phi(0) + b_1 u(1)^{q-1}\phi(1) \right) + \lambda \int_I (mu + au^{p-1})\phi, \quad \forall \phi \in H^1(I).$$

We prove here the following:

Proposition B.1. *Let $b_0, b_1 \in \mathbb{R}$ be arbitrary. Then any nontrivial non-negative weak solution u of* (B.1) *satisfies $u > 0$ in $\overline{I}$.*

Proof. If u is a nonnegative weak solution of (B.1) then, thanks to the inclusion $H^1(I) \subset \mathcal{C}(\overline{I})$ (see [7]) we have $u \in \mathcal{C}(\overline{I})$. Moreover, we claim that $u \in H^2(I)$, so that $u \in \mathcal{C}^1(\overline{I})$. Indeed, from the definition we derive

$$\int_I u'\phi' = \lambda \int_I (mu + au^{p-1})\phi, \quad \forall \phi \in \mathcal{C}_c^1(I).$$

This implies that $(u')' = -au^{p-1}$ in I in the distribution sense. By the chain rule we obtain $mu + au^{p-1} \in H^1(I)$, since $m, a \in \mathcal{C}^1(\overline{I})$. By definition we infer that $u \in H^2(I)$. From the inclusion $H^2(I) \subset \mathcal{C}^1(\overline{I})$, it follows that $u \in \mathcal{C}^1(\overline{I})$.

In fact, by a bootstrap argument and elliptic regularity, we have $u \in \mathcal{C}^2(I)$. Hence, it follows that $u \in \mathcal{C}^1(\overline{I}) \cap \mathcal{C}^2(I)$, and we infer that $u > 0$ in I by the strong maximum principle. In order to show that $u(0) > 0$, we assume by contradiction that $u(0) = 0$. Then the boundary point lemma tells us that $-u'(0) < 0$. However, the boundary condition in (B.1) is understood in the classical sense under the condition $u \in \mathcal{C}^1(\overline{I}) \cap \mathcal{C}^2(I)$, and thus, $u'(0) = 0$, which is a contradiction. Likewise we can show that $u(1) > 0$. □

Remark B.2. Using the same argument as in Proposition B.1, we infer that in the case $N = 1$ nontrivial non-negative solutions of (1.12) satisfy $w_0 > 0$ on $\overline{\Omega}$.

References

[1] S. Agmon, A. Douglis, and L. Nirenberg, Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. I, Comm. Pure Appl. Math. **12**, (1959), 623–727.

[2] S. Alama, G. Tarantello, On semilinear elliptic equations with indefinite nonlinearities, Calc. Var. Partial Differential Equations **1** (1993), 439–475.

[3] A. Ambrosetti, H. Brezis, and G. Cerami, Combined effects of concave and convex nonlinearities in some elliptic problems, J. Funct. Anal. **122**, (1994), 519–543.

[4] C. Bandle, C, M. A. Pozio, A. Tesei, Existence and uniqueness of solutions of nonlinear Neumann problems, Math. Z. **199**, (1988), 257–278.

[5] H. Berestycki, I. Capuzzo-Dolcetta, and L. Nirenberg, Superlinear indefinite elliptic problems and nonlinear Liouville theorems, Topol. Methods Nonlinear Anal. **4**, (1994), 59–78.

[6] H. Berestycki, I. Capuzzo-Dolcetta, and L. Nirenberg, Variational methods for indefinite superlinear homogeneous elliptic problems, NoDEA Nonlinear Differential Equations Appl. **2**, (1995), 553–572.

[7] H. Brezis, Functional analysis, Sobolev spaces and partial differential equations, Universitext. Springer, New York, 2011.

[8] K. J. Brown, Local and global bifurcation results for a semilinear boundary value problem, J. Differential Equations **239**, (2007), 296–310.

[9] K. J. Brown and P. Hess, Stability and uniqueness of positive solutions for a semi-linear elliptic boundary value problem, Differential Integral Equations **3**, (1990), 201–207.

[10] K.J. Brown, S.S. Lin, On the existence of positive eigenfunctions for an eigenvalue problem with indefinite weight function, J. Math. Anal. Appl. **75**, (1980), 112–120.

[11] K. J. Brown and Y. Zhang, The Nehari manifold for a semilinear elliptic equation with a sign-changing weight function, J. Differential Equations **193**, (2003), 481–499.

[12] R. S. Cantrell and C. Cosner, Spatial ecology via reaction-diffusion equations, Wiley Series in Mathematical and Computational Biology, John Wiley & Sons, Ltd., Chichester, 2003.

[13] M. Chipot, M. Fila and P. Quittner, Stationary solutions, blow up and convergence to stationary solutions for semilinear parabolic equations with nonlinear boundary conditions, Acta. Math. Univ. Comenianae **60**, (1991), 35–103.

[14] W. H. Fleming, A selection-migration model in population genetics, J. Math. Biol. **2**, (1975), 219–233.

[15] J. García-Melián, C. Morales-Rodrigo, J. D. Rossi, and A. Suárez, Nonnegative solutions to an elliptic problem with nonlinear absorption and a nonlinear incoming flux on the boundary, Ann. Mat. Pura Appl. (4) **187**, (2008), 459–486.

[16] J. García-Melián, J. D. Rossi, and J. C. Sabina de Lis, Existence and uniqueness of positive solutions to elliptic problems with sublinear mixed boundary conditions, Commun. Contemp. Math. **11**, (2009), 585–613.

[17] D. Gilbarg and N. S. Trudinger, Elliptic partial differential equations of second order, Second edition, Springer-Verlag, Berlin, 1983.

[18] R. Gómez-Reñasco and J. López-Gómez, The effect of varying coefficients on the dynamics of a class of superlinear indefinite reaction-diffusion equations, J. Differential Equations **167**, (2000), 36–72.

[19] J. López-Gómez, On the existence of positive solutions for some indefinite superlinear elliptic problems, Comm. Partial Differential Equations **22**, (1997), 1787–1804.

[20] J. López-Gómez, V. Márquez, and N. Wolanski, Dynamic behavior of positive solutions to reaction-diffusion problems with nonlinear absorption through the boundary, Rev. Un. Mat. Argentina **38**, (1993), 196–209.

[21] C. Morales-Rodrigo and A. Suárez, Some elliptic problems with nonlinear boundary conditions, Spectral theory and nonlinear analysis with applications to spatial ecology, 175–199, World Sci. Publ., Hackensack, NJ, 2005.

[22] T. Ouyang, On the positive solutions of semilinear equations $\Delta u+\lambda u+hu^p = 0$ on compact manifolds. II, Indiana Univ. Math. J. **40**, (1991), 1083–1141.

[23] H. Ramos Quoirin and K. Umezu, The effects of indefinite nonlinear boundary conditions on the structure of the positive solutions set of a logistic equation, J. Differential Equations **257**, (2014), 3935–3977.

[24] H. Ramos Quoirin and K. Umezu, Bifurcation for a logistic elliptic equation with nonlinear boundary conditions: A limiting case, J. Math. Anal. Appl. **428**, (2015), 1265–1285.

[25] J. D. Rossi, Elliptic problems with nonlinear boundary conditions and the Sobolev trace theorem, Stationary partial differential equations, Vol.II, 311–406, Handb. Differ. Equ., Elsevier/North-Holland, Amsterdam, 2005.

[26] G. Stampacchia, Problemi al contorno ellitici, con dati discontinui, dotati di soluzionie hölderiane, Ann. Mat. Pura Appl. (4) **51**, (1960), 1–37.

[27] Nikos M. Stavrakakis, Global bifurcation results for semilinear elliptic equations on $\mathbf{R}^N$: the Fredholm case, J. Differential Equations **142**, (1998), 97–122.

[28] K. Umezu, Bifurcation approach to a logistic elliptic equation with a homogeneous incoming flux boundary condition, J. Differential Equations **252**, (2012), 1146–1168.

[29] K. Umezu, Global structure of supercritical bifurcation with turning points for the logistic elliptic equation with nonlinear boundary conditions, Nonlinear Anal. **89**, (2013), 250–266.

[30] J. L. Vázquez, A strong maximum principle for some quasilinear elliptic equations. Appl. Math. Optim. **12**, (1984), 191–202.

[31] T.-F. Wu, A semilinear elliptic problem involving nonlinear boundary condition and sign-changing potential, Electron. J. Differential Equations **2006**, No.131, 15 pp.

H. Ramos Quoirin
Universidad de Santiago de Chile, Casilla 307, Correo 2, Santiago, Chile

E-mail address: `humberto.ramos@usach.cl`

K. Umezu
Department of Mathematics, Faculty of Education, Ibaraki University, Mito 310-8512, Japan

E-mail address: `kenichiro.umezu.math@vc.ibaraki.ac.jp`